






SUGAR-SUBSTITUTED 2-AZETIDINONES USEFUL AS HYPOCHOLESTEROLEMIC AGENTS**Patent number:** JP10512592T**Publication date:** 1998-12-02**Inventor:****Applicant:****Classification:****- international:** **C07H15/26; C07H15/00;** (IPC1-7): C07H15/203;
A61K31/70**- european:** C07H15/26**Application number:** JP19960517386T 19961029**Priority number(s):** WO1996US16823 19961029; US19950008185P
19951031; US19950570847 19951212**Also published as:** WO9716455 (A1)
 EP0877750 (A1)
 JP2001048895 (A)
 BR9611401 (A)
 EP0877750 (B1)

more >>

Report a data error here

Abstract not available for JP10512592T

Abstract of corresponding document: **WO9716455**

Hypocholesterolemic sugar-substituted 2-azetidinones are disclosed, as well as a method of lowering cholesterol by administering said compounds, pharmaceutical compositions containing them, and the combination of a sugar-substituted 2-azetidinone cholesterol-lowering agent and a cholesterol biosynthesis inhibitor for the treatment and prevention of atherosclerosis.

Data supplied from the **esp@cenet** database - Worldwide

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07H 3/04, 17/02, A61K 31/70	A1	(11) International Publication Number: WO 97/16455 (43) International Publication Date: 9 May 1997 (09.05.97)
(21) International Application Number: PCT/US96/16823 (22) International Filing Date: 29 October 1996 (29.10.96) (30) Priority Data: 60/008,185 31 October 1995 (31.10.95) US 08/570,847 12 December 1995 (12.12.95) US (71) Applicant: SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US). (72) Inventors: YUMIBE, Nathan, P.; 23 South Demarest Avenue, Bergenfield, NJ 07621 (US). ALTON, Kevin, B.; 70 Summit Avenue, Cedar Knolls, NJ 07927 (US). VAN HEEK, Margaret; 1851 Raritan Road, Scotch Plains, NJ 07076 (US). DAVIS, Harry, R.; 64 River Bend Road, Berkeley Heights, NJ 07922 (US). VACCARO, Wayne, D.; 1706 Westover Road, Yardley, PA 19067 (US). (74) Agents: MAGATTI, Anita, W. et al.; Schering-Plough Corporation, Patent Dept. K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: SUGAR-SUBSTITUTED 2-AZETIDINONES USEFUL AS HYPOCHOLESTEROLEMIC AGENTS (57) Abstract Hypocholesterolemic sugar-substituted 2-azetidinones are disclosed, as well as a method of lowering cholesterol by administering said compounds, pharmaceutical compositions containing them, and the combination of a sugar-substituted 2-azetidinone cholesterol-lowering agent and a cholesterol biosynthesis inhibitor for the treatment and prevention of atherosclerosis.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

5

10 **SUGAR-SUBSTITUTED 2-AZETIDINONES**
 USEFUL AS HYPOCHOLESTEROLEMIC AGENTS

BACKGROUND OF THE INVENTION

15 The present invention relates to sugar-substituted 2-azetidinones useful as hypocholesterolemic agents in the treatment and prevention of atherosclerosis, and to the combination of a sugar-substituted 2-azetidinone of this invention and a cholesterol biosynthesis inhibitor for the treatment and prevention of atherosclerosis.

20 Atherosclerotic coronary heart disease represents the major cause for death and cardiovascular morbidity in the western world. Risk factors for atherosclerotic coronary heart disease include hypertension, diabetes mellitus, family history, male gender, cigarette smoke and serum cholesterol. A total cholesterol level in excess of 225-250 mg/dl is associated with significant elevation of risk.

25 Cholesteryl esters are a major component of atherosclerotic lesions and the major storage form of cholesterol in arterial wall cells. Formation of cholesteryl esters is also a key step in the intestinal absorption of dietary cholesterol. In addition to regulation of dietary cholesterol, the regulation of whole-body cholesterol homeostasis in humans and animals involves modulation of cholesterol biosynthesis, bile acid biosynthesis, and the catabolism of the cholesterol-containing plasma lipoproteins. The liver is the major organ responsible for cholesterol biosynthesis and catabolism and, for this reason, it is a prime determinant of plasma cholesterol levels. The liver
30 is the site of synthesis and secretion of very low density lipoproteins (VLDL) which are subsequently metabolized to low density lipoproteins
35

-2-

(LDL) in the circulation. LDL are the predominant cholesterol-carrying lipoproteins in the plasma and an increase in their concentration is correlated with increased atherosclerosis.

When cholesterol absorption in the intestines is reduced,
5 by whatever means, less cholesterol is delivered to the liver. The consequence of this action is a decreased hepatic lipoprotein (VLDL) production and an increase in the hepatic clearance of plasma cholesterol, mostly as LDL. Thus, the net effect of an inhibition of intestinal cholesterol absorption is a decrease in plasma cholesterol
10 levels.

Several 2-azetidinone compounds have been reported as being useful in lowering cholesterol and/or in inhibiting the formation of cholesterol-containing lesions in mammalian arterial walls: WO 93/02048 describes 2-azetidinone compounds wherein the 3-position
15 substituent is arylalkylene, arylalkenylene or arylalkylene wherein the alkylene, alkenylene or alkyleneportion is interrupted by a hetero atom, phenylene or cycloalkylene; WO 94/17038 describes 2-azetidinone compounds wherein the 3-position substituent is an arylalkylspirocyclic group; WO 95/08532 describes 2-azetidinone compounds wherein the
20 3-position substituent is an arylalkylene group substituted in the alkylene portion by a hydroxy group; PCT/US95/03196, filed March 22, 1995, describes compounds wherein the 3-position substituent is an aryl(oxo or thio)alkylene group substituted in the alkylene portion by a hydroxy group; and U.S. Serial No. 08/463,619, filed June 5, 1995,
25 describes the preparation of compounds wherein the 3-position substituent is an arylalkylene group substituted in the alkylene portion by a hydroxy group, and wherein the alkylene group is attached the the azetidinone ring by a -S(O)₀₋₂- group. The cited patent applications are incorporated herein by reference.

30 Also, European Patent 199,630 and European Patent Application 337,549 disclose elastase inhibitory substituted azetidinones said to be useful in treating inflammatory conditions resulting in tissue destruction which are associated with various disease states, e.g. atherosclerosis.

35 Other known hypocholesterolemics include plant extracts such as sapogenins, in particular tigogenin and diosgenin. Glycoside

-3-


derivatives of tigogenin and/or diosgenin are disclosed in PCT International publications WO 94/00480 and WO 95/18143.

The inhibition of cholesterol biosynthesis by 3-hydroxy-3-methylglutaryl coenzyme A reductase (EC1.1.1.34) inhibitors has been shown to be an effective way to reduce plasma cholesterol (Witzum, *Circulation*, 80, 5 (1989), p. 1101-1114) and reduce atherosclerosis. Combination therapy of an HMG CoA reductase inhibitor and a bile acid sequestrant has been demonstrated to be more effective in human hyperlipidemic patients than either agent in monotherapy (Illingworth, *Drugs*, 36 (Suppl. 3) (1988), p. 63-71).

SUMMARY OF THE INVENTION

The present invention relates to sugar-substituted 2-azetidinones, especially to glucose-derived conjugates of cholesterol-lowering 2-azetidinones having an aryl or substituted aryl group as a substituent at the 1-position and having a hydroxy-substituted phenyl group, especially a 4-hydroxyphenyl group, at the 4-position.



Compounds of the present invention are represented by the formula I



or a pharmaceutically acceptable salt thereof, wherein

R²⁶ is H or OG¹;

G and G¹ are independently selected from the group consisting of



provided that when R²⁶ is H or

-4-

OH, G is not H;

R, R^a and R^b are independently selected from the group consisting of H, -OH, halogeno, -NH₂, azido, (C₁-C₆)alkoxy(C₁-C₆)-alkoxy or -W-R³⁰;

5 W is independently selected from the group consisting of -NH-C(O)-, -O-C(O)-, -O-C(O)-N(R³¹)-, -NH-C(O)-N(R³¹)- and -O-C(S)-N(R³¹)-;

R² and R⁶ are independently selected from the group consisting of H, (C₁-C₆)alkyl, aryl and aryl(C₁-C₆)alkyl;

10 R³, R⁴, R⁵, R⁷, R^{3a} and R^{4a} are independently selected from the group consisting of H, (C₁-C₆)alkyl, aryl(C₁-C₆)alkyl, -C(O)(C₁-C₆)alkyl and -C(O)aryl;

R³⁰ is independently selected from the group consisting of R³²-substituted T, R³²-substituted-T-(C₁-C₆)alkyl, R³²-substituted-(C₂-C₄)alkenyl, R³²-substituted-(C₁-C₆)alkyl, R³²-substituted-(C₃-C₇)cycloalkyl and R³²-substituted-(C₃-C₇)cycloalkyl(C₁-C₆)alkyl;

R³¹ is independently selected from the group consisting of H and (C₁-C₄)alkyl;


20 T is independently selected from the group consisting of phenyl, furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl and pyridyl;

R³² is independently selected from 1-3 substituents independently selected from the group consisting of halogeno, (C₁-C₄)alkyl, -OH, phenoxy, -CF₃, -NO₂, (C₁-C₄)alkoxy, methylenedioxy, 25 oxo, (C₁-C₄)alkylsulfanyl, (C₁-C₄)alkylsulfinyl, (C₁-C₄)alkylsulfonyl, -N(CH₃)₂, -C(O)-NH(C₁-C₄)alkyl, -C(O)-N((C₁-C₄)alkyl)₂, -C(O)-(C₁-C₄)alkyl, -C(O)-(C₁-C₄)alkoxy and pyrrolidinylcarbonyl; or R³² is a covalent bond and R³¹, the nitrogen to which it is attached and R³² form a pyrrolidinyl, piperidinyl, N-methyl-piperazinyl, indolyl or morpholyl group, or a (C₁-C₄)alkoxycarbonyl-substituted pyrrolidinyl, piperidinyl, 30 N-methylpiperazinyl, indolyl or morpholyl group;

Ar¹ is aryl or R¹⁰-substituted aryl;

Ar² is aryl or R¹¹-substituted aryl;

Q is a bond or, with the 3-position ring carbon of the azetidinone,



R¹ is selected from the group consisting of


-(CH₂)_q-, wherein q is 2-6, provided that when Q forms a spiro ring, q can also be zero or 1;

5 $-(\text{CH}_2)_e\text{-E-(CH}_2)_r$, wherein E is -O-, -C(O)-, phenylene, -NR²²- or -S(O)₀₋₂-, e is 0-5 and r is 0-5, provided that the sum of e and r is 1-6;

-(C₂-C₆)alkenylene-; and

10 $-(CH_2)_f-V-(CH_2)_g-$, wherein V is C₃-C₆ cycloalkylene, f is 1-5 and g is 0-5, provided that the sum of f and g is 1-6;


R¹² is



R¹³ and R¹⁴ are independently selected from the group consisting of -CH₂-, -CH(C₁-C₆ alkyl)-, -C(di-(C₁-C₆) alkyl)-, -CH=CH- and -C(C₁-C₆ alkyl)=CH-; or R¹² together with an adjacent R¹³, or R¹² together with an adjacent R¹⁴, form a -CH=CH- or a -CH=C(C₁-C₆ alkyl)- group;

a and b are independently 0, 1, 2 or 3, provided both are not zero;
provided that when R¹³ is -CH=CH- or -C(C₁-C₆ alkyl)=CH-, a is 1;
20 provided that when R¹⁴ is -CH=CH- or -C(C₁-C₆ alkyl)=CH-, b is 1;
provided that when a is 2 or 3, the R¹³'s can be the same or different;
and provided that when b is 2 or 3, the R¹⁴'s can be the same or
different;

and when Q is a bond, R¹ also can be:



M is -O-, -S-, -S(O)- or -S(O)₂-;

X, Y and Z are independently selected from the group consisting of -CH₂-, -CH(C₁-C₆)alkyl- and -C(di-(C₁-C₆)alkyl);

R¹⁰ and R¹¹ are independently selected from the group
30 consisting of 1-3 substituents independently selected from the group

-6-

- consisting of (C₁-C₆)alkyl, -OR¹⁹, -O(CO)R¹⁹, -O(CO)OR²¹,
 -O(CH₂)₁₋₅OR¹⁹, -O(CO)NR¹⁹R²⁰, -NR¹⁹R²⁰, -NR¹⁹(CO)R²⁰,
 -NR¹⁹(CO)OR²¹, -NR¹⁹(CO)NR²⁰R²⁵, -NR¹⁹SO₂R²¹, -COOR¹⁹,
 -CONR¹⁹R²⁰, -COR¹⁹, -SO₂NR¹⁹R²⁰, S(O)₀₋₂R²¹,
 5 -O(CH₂)₁₋₁₀-COOR¹⁹, -O(CH₂)₁₋₁₀CONR¹⁹R²⁰, -(C₁-C₆ alkylene)-
 COOR¹⁹, -CH=CH-COOR¹⁹, -CF₃, -CN, -NO₂ and halogen;
 R¹⁵ and R¹⁷ are independently selected from the group
 consisting of -OR¹⁹, -O(CO)R¹⁹, -O(CO)OR²¹ and -O(CO)NR¹⁹R²⁰; R¹⁶
 and R¹⁸ are independently selected from the group consisting of H,
 10 (C₁-C₆)alkyl and aryl; or R¹⁵ and R¹⁶ together are =O, or R¹⁷ and R¹⁸
 together are =O;
 d is 1, 2 or 3;
 h is 0, 1, 2, 3 or 4;
 s is 0 or 1; t is 0 or 1; m, n and p are independently 0-4; provided
 15 that at least one of s and t is 1, and the sum of m, n, p, s and t is 1-6;
 provided that when p is 0 and t is 1, the sum of m, s and n is 1-5; and
 provided that when p is 0 and s is 1, the sum of m, t and n is 1-5;
 v is 0 or 1;
 j and k are independently 1-5, provided that the sum of j, k and v
 20 is 1-5;
- $$\begin{array}{c} \text{R}^{15} \\ | \\ -\text{X}_j - (\text{C})_v - \text{Y}_k - \text{S}(\text{O})_{0-2} - \\ | \\ \text{R}^{16} \end{array}$$
- and when Q is a bond and R¹ is $\begin{array}{c} \text{R}^{15} \\ | \\ -\text{X}_j - (\text{C})_v - \text{Y}_k - \text{S}(\text{O})_{0-2} - \\ | \\ \text{R}^{16} \end{array}$, Ar¹ can
 also be pyridyl, isoxazolyl, furanyl, pyrrolyl, thienyl, imidazolyl, pyrazolyl,
 thiazolyl, pyrazinyl, pyrimidinyl or pyridazinyl;
 R¹⁹ and R²⁰ are independently selected from the group
 25 consisting of H, (C₁-C₆)alkyl, aryl and aryl-substituted (C₁-C₆)alkyl;
 R²¹ is (C₁-C₆)alkyl, aryl or R²⁴-substituted aryl;
 R²² is H, (C₁-C₆)alkyl, aryl (C₁-C₆)alkyl, -C(O)R¹⁹ or -COOR¹⁹;
 R²³ and R²⁴ are independently 1-3 groups independently
 selected from the group consisting of H, (C₁-C₆)alkyl, (C₁-C₆)alkoxy,
 30 -COOH, NO₂, -NR¹⁹R²⁰, -OH and halogeno; and
 R²⁵ is H, -OH or (C₁-C₆)alkoxy.

-7-

Ar² is preferably phenyl or R¹¹-phenyl, especially (4-R¹¹)-substituted phenyl. Preferred definitions of R¹¹ are lower alkoxy, especially methoxy, and halogeno, especially fluoro.

Ar¹ is preferably phenyl or R¹⁰-substituted phenyl, especially (4-R¹⁰)-substituted phenyl. A preferred definition of R¹⁰ is halogeno, especially fluoro.

There are several preferred definitions for the -R¹-Q-combination of variables:

Q is a bond and R¹ is lower alkylene, preferably propylene;

Q is a spiro group as defined above, wherein preferably R¹³ and R¹⁴ are each ethylene and R¹² is $\begin{smallmatrix} | \\ -\text{CH}- \\ | \end{smallmatrix}$ or $\begin{smallmatrix} | \\ -\text{C}(\text{OH})- \\ | \end{smallmatrix}$, and R¹ is $-(\text{CH}_2)_q$ wherein q is 0-6;

Q is a bond and R¹ is $-\text{M}-\text{Y}_d-\overset{\overset{\text{R}^{15}}{|}}{\underset{\underset{\text{R}^{16}}{|}}{\text{C}}}-\text{Z}_h-$ wherein the variables are chosen such that R¹ is -O-CH₂-CH(OH)-;

Q is a bond and R¹ is $-\text{X}_m-\overset{\overset{\text{R}^{17}}{|}}{\underset{\underset{\text{R}^{18}}{|}}{\text{C}}}_s-\text{Y}_n-\overset{\overset{\text{R}^{15}}{|}}{\underset{\underset{\text{R}^{16}}{|}}{\text{C}}}_t-\text{Z}_p-$ wherein the variables are chosen such that R¹ is -CH(OH)-(CH₂)₂-; and


Q is a bond and R¹ is $-\text{X}_j-\overset{\overset{\text{R}^{15}}{|}}{\underset{\underset{\text{R}^{16}}{|}}{\text{C}}}_v-\text{Y}_k-\text{S}(\text{O})_{0-2}-$ wherein the variables are chosen such that R¹ is -CH(OH)-CH₂-S(O)₀₋₂-.

A preferred compound of formula I, therefore, is one wherein G and G¹ are as defined above and in which the remaining variables have the following definitions:


Ar¹ is phenyl or R¹⁰-substituted phenyl, wherein R¹⁰ is halogeno;

Ar² is phenyl or R¹¹-phenyl, wherein R¹¹ is 1 to 3 substituents independently selected from the group consisting of C₁-C₆ alkoxy and halogeno;

Q is a bond and R¹ is lower alkylene; Q, with the 3-position

ring carbon of the azetidinone, forms the group $(R^{14})_b$  wherein preferably R¹³ and R¹⁴ are each ethylene and a and b are each 1, and wherein R¹² is $\text{-}\overset{|}{\text{C}}\text{H-}$ or $\text{-}\overset{|}{\text{C}}(\text{OH})\text{-}$; Q is a bond and R¹ is $\text{-O-CH}_2\text{-CH(OH)-}$; Q is a bond and R¹ is $\text{-CH(OH)-(CH}_2\text{)}_2\text{-}$; or Q is a bond and R¹ is $\text{-CH(OH)-CH}_2\text{-S(O)}_{0-2}\text{-}$.


Preferred variables for G and G¹ groups of the formula



are as follows:

10 R², R³, R⁴, R⁵, R⁶ and R⁷ are independently selected from the group consisting of H, (C₁-C₆)alkyl, benzyl and acetyl.

Preferred variables for group G or G¹ of the formula



are as follows:

R³, R^{3a}, R⁴ and R^{4a} are selected from the group consisting of H,
15 (C₁-C₆)alkyl, benzyl and acetyl;


R, R^a and R^b are independently selected from the group consisting of H, -OH, halogeno, -NH₂, azido, (C₁-C₆)alkoxy(C₁-C₆)alkoxy and -W-R³⁰, wherein W is -O-C(O)- or -O-C(O)-NR³¹-, R³¹ is H and R³⁰ is (C₁-C₆)alkyl, -C(O)-(C₁-C₄)alkoxy-(C₁-C₆)alkyl, T, T-(C₁-C₆)alkyl, or T or T-(C₁-C₆)alkyl wherein T is substituted by one or two halogeno or (C₁-C₆)alkyl groups.

Preferred R³⁰ substituents are 2-fluorophenyl, 2,4-difluorophenyl, 2,6-dichlorophenyl, 2-methylphenyl, 2-thienylmethyl, 2-methoxycarbonylethyl, thiazol-2-yl-methyl, 2-furyl, 2-methoxycarbonylbutyl and phenyl. Preferred combinations of R, R^a and R^b are as follows: 1) R, R^a and R^b are independently -OH or -O-C(O)-NH-R³⁰, especially wherein

-9-


- R^a is -OH and R and R^b are -O-C(O)-NH- R^{30} and R^{30} is selected from the preferred substituents identified above, or wherein R and R^a are -OH and R^b is -O-C(O)-NH- R^{30} wherein R^{30} is 2-fluorophenyl, 2,4-difluorophenyl, 2,6-dichlorophenyl; 2) R^a is -OH, halogeno, azido or (C₁-C₆)-alkoxy(C₁-C₆)alkoxy, R^b is H, halogeno, azido or (C₁-C₆)alkoxy(C₁-C₆)-alkoxy, and R is -O-C(O)-NH- R^{30} , especially compounds wherein R^a is -OH, R^b is H and R^{30} is 2-fluorophenyl; 3) R, R^a and R^b are independently -OH or -O-C(O)- R^{30} and R^{30} is (C₁-C₆)alkyl, T, or T substituted by one or two halogeno or (C₁-C₆)alkyl groups, especially compounds wherein R is -OH and R^a and R^b are -O-C(O)- R^{30} wherein R^{30} is 2-furyl; and 4) R, R^a and R^b are independently -OH or halogeno. Three additional classes of preferred are compounds are those wherein the C1' anomeric oxy is beta, wherein the C2' anomeric oxy is beta, and wherein the R group is alpha.

- 15 G and G¹ are preferably selected from:



20

-10-



wherein Ac is acetyl and Ph is phenyl. Preferably, R²⁶ is H or OH, more preferably H. The -O-G substituent is preferably in the 4-position of the phenyl ring to which it is attached.

5 This invention also relates to the use of a sugar-substituted 2-azetidinone, especially one of formula I, as a hypocholesterolemic agent in a mammal in need of such treatment.

 In another aspect, the invention relates to a pharmaceutical composition comprising a sugar-substituted 2-azetidinone, especially
10 one of formula I, in a pharmaceutically acceptable carrier.

 The present invention also relates to a method of reducing hepatic cholesterol ester levels, a method of reducing plasma cholesterol levels, and to a method of treating or preventing atherosclerosis, comprising administering to a mammal in need of such
15 treatment an effective amount of a combination of a sugar-substituted 2-azetidinone of this invention, especially one of formula I, and a cholesterol biosynthesis inhibitor. That is, the present invention relates to the use of a sugar-substituted 2-azetidinone for combined use with a cholesterol biosynthesis inhibitor (and, similarly, use of a cholesterol
20 biosynthesis inhibitor for combined use with a sugar-substituted 2-azetidinone) to treat or prevent atherosclerosis or to reduce plasma cholesterol levels.

 In yet another aspect, the invention relates to a pharmaceutical composition comprising an effective amount of a sugar-
25 substituted 2-azetidinone, a cholesterol biosynthesis inhibitor, and a pharmaceutically acceptable carrier. In a final aspect, the invention relates to a kit comprising in one container an effective amount of a sugar-substituted 2-azetidinone in a pharmaceutically acceptable carrier, and in a separate container, an effective amount of a cholesterol
30 biosynthesis inhibitor in a pharmaceutically acceptable carrier.

DETAILED DESCRIPTION:

As used herein, the term "alkyl" or "lower alkyl" means straight or branched alkyl chains of 1 to 6 carbon atoms and "alkoxy" similarly refers to alkoxy groups having 1 to 6 carbon atoms.

5 "Alkenyl" means straight or branched carbon chains having one or more double bonds in the chain, conjugated or unconjugated. Similarly, "alkynyl" means straight or branched carbon chains having one or more triple bonds in the chain. Where an alkyl, alkenyl or alkynyl chain joins two other variables and is therefore bivalent, the terms
10 alkylene, alkenylene and alkynylene are used.

"Cycloalkyl" means a saturated carbon ring of 3 to 6 carbon atoms, while "cycloalkylene" refers to a corresponding bivalent ring, wherein the points of attachment to other groups include all positional isomers.

15 "Halogeno" refers to fluorine, chlorine, bromine or iodine radicals.

"Aryl" means phenyl, naphthyl, indenyl, tetrahydronaphthyl or indanyl. "Phenylene" means a bivalent phenyl group, including ortho, meta and para-substitution. R²⁴-benzyl and R²⁴-benzyloxy refer to
20 benzyl and benzyloxy radicals which are substituted on the phenyl ring.

The above statements, wherein, for example, R¹⁹, R²⁰ and R²⁵ are said to be independently selected from a group of substituents, means that R¹⁹, R²⁰ and R²⁵ are independently selected, but also that where an R¹⁹, R²⁰ or R²⁵ variable occurs more than once in a molecule,
25 those occurrences are independently selected (e.g., if R¹⁰ is -OR¹⁹ wherein R¹⁹ is hydrogen, R¹¹ can be -OR¹⁹ wherein R¹⁹ is lower alkyl). Those skilled in the art will recognize that the size and nature of the substituent(s) will affect the number of substituents which can be present.

30 Compounds of the invention have at least one asymmetrical carbon atom and therefore all isomers, including diastereomers and rotational isomers are contemplated as being part of this invention. The invention includes d and l isomers in both pure form and in admixture, including racemic mixtures. Isomers can be prepared
35 using conventional techniques, either by reacting optically pure or

optically enriched starting materials or by separating isomers of a compound of formula I.

Those skilled in the art will appreciate that for some compounds of formula I, one isomer will show greater pharmacological activity than other isomers.

Compounds of the invention with an amino group can form pharmaceutically acceptable salts with organic and inorganic acids. Examples of suitable acids for salt formation are hydrochloric, sulfuric, phosphoric, acetic, citric, oxalic, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and carboxylic acids well known to those in the art. The salt is prepared by contacting the free base form with a sufficient amount of the desired acid to produce a salt. The free base form may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous sodium bicarbonate. The free base form differs from its respective salt form somewhat in certain physical properties, such as solubility in polar solvents, but the salt is otherwise equivalent to its respective free base forms for purposes of the invention.


Certain compounds of the invention are acidic (e.g., those compounds which possess a carboxyl group). These compounds form pharmaceutically acceptable salts with inorganic and organic bases. Examples of such salts are the sodium, potassium, calcium, aluminum, gold and silver salts. Also included are salts formed with pharmaceutically acceptable amines such as ammonia, alkyl amines, hydroxyalkylamines, N-methylglucamine and the like.

Cholesterol biosynthesis inhibitors for use in the combination of the present invention include HMG CoA reductase inhibitors such as lovastatin, pravastatin, fluvastatin, simvastatin and CI-981; HMG CoA synthetase inhibitors, for example L-659,699 ((E,E-11-[3'R-(hydroxy-methyl)-4'-oxo-2'R-oxetanyl]-3,5,7R-trimethyl-2,4-undecadienoic acid); squalene synthesis inhibitors, for example squalastatin 1; and squalene epoxidase inhibitors, for example, NB-598 ((E)-N-ethyl-N-(6,6-dimethyl-2-hepten-4-ynyl)-3-[(3,3'-bithiophen-5-yl)methoxy]benzene-methanamine hydrochloride). Preferred HMG CoA reductase inhibitors are lovastatin, pravastatin, fluvastatin and simvastatin.


-13-

The cholesterol-lowering 2-azetidinone portions of the compounds of formula I can be prepared by known methods, for example WO 93/02048 describes the preparation of compounds wherein -R¹-Q- is alkylene, alkenylene or alkylene interrupted by a hetero atom, phenylene or cycloalkylene; WO 94/17038 describes the preparation of compounds wherein Q is a spirocyclic group; WO 95/08532 describes the preparation of compounds wherein -R¹-Q- is a hydroxy-substituted alkylene group; PCT/US95/03196 describes compounds wherein -R¹-Q- is a hydroxy-substituted alkylene attached to the Ar¹ moiety through an -O- or S(O)₀₋₂- group; and U.S. Serial No. 08/463,619, filed June 5, 1995, describes the preparation of compounds wherein -R¹-Q- is a hydroxy-substituted alkylene group attached the the azetidinone ring by a -S(O)₀₋₂- group.


Compounds of the present invention are generally prepared by reacting a 4-(hydroxy- or dihydroxy)-phenyl-2-azetidinone with a sugar derivative. For example, an azetidinone of formula II, wherein R^{26A} is H or OH, is reacted with one equivalent of a sugar derivative of formula III:




wherein R³⁰ is hydrogen or -CNHCCl₃ and the remaining variables are as defined above to obtain a compound of formula IA, wherein R^{26A} is H or OH. To prepare a compound of formula IB, wherein R²⁶ is OG¹, wherein G¹ is not H, and G is H, an azetidinone of formula IIA, wherein R²⁶ is OH and R²⁷ is a suitable hydroxy protecting group, is reacted with a sugar derivative of formula IIIA, wherein R³⁰ is as defined above, followed by removal of the R²⁷ protecting group:



To prepare a compound of formula IC, wherein both G¹ and G are the same, but are not H, a dihydroxy compound of formula IIC is reacted with an excess of G-OR³⁰:



- 5 To prepare compounds of formula ID wherein G and G¹ are both not H and are not the same sugar derivative, a compound of formula IA wherein R^{26A} is OH can be reacted with a sugar of the formula G¹-OR³⁰. Alternatively, one of the hydroxy substituents on the 4-position phenyl of a compound of formula IIC is protected prior to reaction with the sugar derivative to be attached to the unprotected hydroxy group, and after reaction with the first sugar derivative, the hydroxy-protecting group is removed and the second sugar derivative is reacted with the previously-protected hydroxy group. For example:




- 15 Sugars and the derivatives thereof as defined by G-OR³⁰ and G¹-OR³⁰ are known in the art or are readily prepared by known methods.

- 20 Preferably, the reactions described above involve a sugar derivative wherein the non-reactive hydroxy groups are protected by suitable protecting groups as defined above for R², R³, R^{3a}, R⁴, R^{4a}, R⁵ and R⁷ other than hydrogen, preferably lower alkyl, acetyl or benzyl, which groups can be removed after the reaction to provide the sugar conjugate. When the 1- and 3-position side chains of the 2-azetidinone include substituent groups which are reactive under the conditions

-15-

used, said reactive groups are protected by suitable protecting groups prior to reaction with the sugar or the derivative thereof, and the protecting groups are subsequently removed. Depending on the nature of the protecting groups, the protecting groups on the sugar portion and on the 1- and 3-position side chains of the azetidinone can be removed sequentially or simultaneously.

For example, compounds of formula I wherein $\text{Ar}^1\text{-R}^1\text{-Q-}$ is $\text{Ar}^1\text{-CH(OH)-}(\text{CH}_2)_2\text{-}$, i.e. compounds of formula Ia and Ib, can be prepared according to the following reaction scheme, wherein an azetidinone of formula IIa is reacted with a sugar derivative of the formula G-OCNHCCl_3 . The scheme is shown for a compound wherein R_{26} is H and a specific G-OCNHCCl_3 group is exemplified, but a similar procedure can be used to prepare compounds wherein R_{26} is $-\text{OG}^1$ and for other G-OCNHCCl_3 groups:



-16-

In the first step, the azetidinone of formula IIa is reacted with the sugar derivative of formula IIIa in the presence of a coupling agent such as BF_3 etherate in an inert solvent such as CH_2Cl_2 . The reaction is carried out at temperatures of -20 to -25°C for a period of about two hours. In the


5 second step, either the sugar-substituted azetidinone of formula IV is treated with a base such as triethylamine in a solvent such as methanol and water to remove the acetyl and alkyl protecting groups to obtain a compound of formula Ia, or the sugar-substituted azetidinone of formula IV is treated with a reagent such as KCN in a solvent such as methanol

10 to remove the acetyl protecting groups but leave the alkyl protecting group to obtain a compound of formula Ib. The compound of formula Ib can be further reduced by a reagent such as LiOH to obtain the compound of formula Ia.


Compounds of formula I wherein $\text{Ar}^1\text{-R}^1\text{-Q-}$ is $\text{Ar}^1\text{-(CH}_2\text{)}_3\text{-}$, i.e. compounds of formula Ic, can be prepared according to the following

15 reaction scheme, wherein an azetidinone of formula IIb is reacted with a sugar derivative of the formula G-OH. The scheme is shown for a compound wherein R^{26} is hydrogen and with a specific G-OH group, but a similar procedure can be used to prepare compounds wherein R^{26} is

20 -OG^1 and for other G-OH groups:



-17-



In the first step, the azetidinone of formula IIb is reacted with a sugar derivative of formula IIIb in an inert solvent such as tetrahydrofuran in the presence of n-tributylphosphine and 1,1'-(azodicarbonyl)dipiperidine.


- 5 The resultant sugar-substituted azetidinone is reduced with a reagent such as $\text{Pd(OH)}_2/\text{C}$ in an alcoholic solvent under H_2 gas to remove the benzyl protecting groups to obtain a compound of formula I.

- Starting materials of formula IIb are known. Compounds of formula IIa can be prepared from the corresponding (3-hydroxy-3 Ar^1 -propyl)-2-azetidinone by treatment with acetic anhydride and dimethylaminopyridine (DMAP) in an inert solvent such as CH_2Cl_2 to obtain the corresponding di-acetyl compound, followed by treatment with guanidine to obtain the 4-hydroxyphenyl compound. Starting materials of formula II wherein $\text{Ar}^1\text{-R}^1\text{-Q-}$ is as defined above for formula I can be prepared by similar methods or others well known in the art.
- 10
- 15

- Starting materials of formula IIIb are known in the art or prepared by well known methods. Compounds of formula IIIa are prepared by treating the corresponding compound of formula IIIb with trichloroacetonitrile in an inert solvent such as CH_2Cl_2 in the presence of Cs_2CO_3 .
- 20

Reactive groups not involved in the above processes can be protected during the reactions with conventional protecting groups which can be removed by standard procedures after the reaction. The following Table 1 shows some typical protecting groups:

Table 1

Group to be Protected	Group to be Protected and Protecting Group
-COOH	-COOalkyl, -COObenzyl, -COOphenyl
>NH	$\begin{array}{l} \diagup \text{NCOalkyl}, \diagup \text{NCObenzyl}, \diagup \text{NCOphenyl} \\ \diagdown \text{NCH}_2\text{OCH}_2\text{CH}_2\text{Si}(\text{CH}_3)_3, \diagdown \text{NC}(\text{O})\text{OC}(\text{CH}_3)_3, \\ \diagdown \text{N-benzyl}, \diagdown \text{NSi}(\text{CH}_3)_3, \diagdown \text{NSi}(\text{CH}_3)_2\text{C}(\text{CH}_3)_2 \end{array}$
-NH ₂	
-OH	$\begin{array}{l} -\text{OCH}_3, -\text{OCH}_2\text{OCH}_3, -\text{OSi}(\text{CH}_3)_3, -\text{OSi}(\text{CH}_3)_2\text{C}(\text{CH}_3)_2 \\ \text{or } -\text{OCH}_2\text{phenyl} \end{array}$

We have found that the compounds of this invention lower plasma lipid levels and hepatic cholesterol ester levels. Compounds of this invention have been found to inhibit the intestinal absorption of cholesterol and to significantly reduce the formation of liver cholesteryl esters in animal models. Thus, compounds of this invention are hypocholesterolemic agents by virtue of their ability to inhibit the esterification and/or intestinal absorption of cholesterol; they are therefore useful in the treatment and prevention of atherosclerosis in mammals, in particular in humans.

Compared to the 2-azetidinone cholesterol lowering agents which are not sugar-substituted, the compounds of this invention have several pharmacological and physical advantages. The compounds are absorbed at a slower rate, give lower plasma levels and higher intestinal levels. Previous testing indicated the intestine as the likely site of activity of the 2-azetidinone compounds lacking a sugar substituent. See E. J. Sybertz et al, "SCH 48461, a Novel Inhibitor of Cholesterol Absorption," *Atherosclerosis X*, ed. F.P. Woodward et al (Elsevier, 1995), pp. 311-315; and B.G. Salisbury et al,

"Hypercholesterolemic Activity of a Novel Inhibitor of Cholesterol Absorption," *Atherosclerosis*, 115 (1995), pp. 45-63. The instantly claimed compounds, which are excreted in the bile, provide efficient delivery of the compound to the desired site while minimizing systemic exposure, thereby decreasing potential toxicity problems.

In addition to the compound aspect, the present invention also relates to a method of lowering plasma cholesterol levels, which method comprises administering to a mammal in need of such treatment a hypocholesterolemic effective amount of a compound of formula I of this invention. The compound is preferably administered in a pharmaceutically acceptable carrier suitable for oral administration.

The present invention also relates to a pharmaceutical composition comprising a compound of formula I of this invention and a pharmaceutically acceptable carrier. The compounds of formula I can be administered in any conventional oral dosage form such as capsules, tablets, powders, cachets, suspensions or solutions. The formulations and pharmaceutical compositions can be prepared using conventional pharmaceutically acceptable excipients and additives and conventional techniques. Such pharmaceutically acceptable excipients and additives include non-toxic compatible fillers, binders, disintegrants, buffers, preservatives, anti-oxidants, lubricants, flavorings, thickeners, coloring agents, emulsifiers and the like.

The daily hypocholesterolemic dose of a compound of formula I is about 0.001 to about 30 mg/kg of body weight per day, preferably about 0.001 to about 1 mg/kg. For an average body weight of 70kg, the dosage level is therefore from about 0.1 to about 100 mg of drug per day, given in a single dose or 2-4 divided doses. The exact dose, however, is determined by the attending clinician and is dependent on the potency of the compound administered, the age, weight, condition and response of the patient.

For the combinations of this invention wherein the substituted azetidinone is administered in combination with a cholesterol biosynthesis inhibitor, the typical daily dose of the cholesterol biosynthesis inhibitor is 0.1 to 80 mg/kg of mammalian weight per day administered in single or divided dosages, usually once or twice a day: for example, for HMG CoA reductase inhibitors, about 10

-20-

- to about 40 mg per dose is given 1 to 2 times a day, giving a total daily dose of about 10 to 80 mg per day, and for the other cholesterol biosynthesis inhibitors, about 1 to 1000 mg per dose is given 1 to 2 times a day, giving a total daily dose of about 1 mg to about 2 g per day.
- 5 The exact dose of any component of the combination to be administered is determined by the attending clinician and is dependent on the potency of the compound administered, the age, weight, condition and response of the patient.

- Where the components of a combination are administered separately, the number of doses of each component given per day may not necessarily be the same, e.g. where one component may have a greater duration of activity, and will therefore need to be administered less frequently.

- Since the present invention relates to the reduction of plasma cholesterol levels by treatment with a combination of active ingredients wherein said active ingredients may be administered separately, the invention also relates to combining separate pharmaceutical compositions in kit form. That is, a kit is contemplated wherein two separate units are combined: a cholesterol biosynthesis inhibitor pharmaceutical composition and a sugar-substituted 2-azetidinone absorption inhibitor pharmaceutical composition. The kit will preferably include directions for the administration of the separate components. The kit form is particularly advantageous when the separate components must be administered in different dosage forms (e.g. oral and parenteral) or are administered at different dosage intervals.

Following are examples of preparing compounds of formula I. The stereochemistry listed is relative stereochemistry unless otherwise noted. The terms cis and trans refer to the relative orientations at the β -lactam 3- and 4-positions.

30

Preparation A

1-(4-Fluorophenyl)-3(R)-[3(S)-acetyloxy-3-(4-fluorophenyl)-propyl]-4(S)-(4-hydroxyoxyphenyl)-2-azetidinone

Step 1: 1-(4-Fluorophenyl)-3(R)-[3(S)-acetyloxy-3-(4-fluorophenyl)-propyl]-4(S)-(4-acetyloxyphenyl)-2-azetidinone

35

Add acetic anhydride (1.03 mL, 10.96 mmol) to a room temperature solution of 1-(4-fluorophenyl)-3(R)-[3(S)-hydroxy-3-(4-

-21-

fluorophenyl)propyl]]-4(S)-(4-hydroxyphenyl)-2-azetidinone (2.04 g, 4.98 mmol) and dimethylaminopyridine (DMAP) (1.46 g, 11.96 mmol) in tetrahydrofuran (THF) (15 mL). After TLC (5% CH₃OH/toluene) indicates consumption of starting material (10 min), dilute the mixture with ether (Et₂O), wash with 1M HCl and brine, dry over anhydrous Na₂SO₄, concentrate to a clear foam 2.47 g (100%) and use without further purification. NMR (400 MHz, CDCl₃): 7.33(2H, d, J=8.6 Hz), 7.27(2H, m), 7.21(2H, m), 7.11(2H, d, J=8.5 Hz), 7.02(2H, t, J=8.6 Hz), 6.94(2H, d, J=8.5 Hz), 5.70(1H, t, J= 7Hz), 4.60(1H, d, J=2.4 Hz), 3.06(1H, dt, J=7.9, 2.4 Hz), 2.31(3H, s), 2.06(3H, s), 2.03(1H, m), 1.86(2H, M). HRMS (FAB): calcd. for M+H: C₂₈H₂₅NO₅F₂, 493.1701; found 493.1695.

Step 2: Add sodium ethoxide (0.338 g, 4.97 mmol) to a room temperature solution of guanadine hydrochloride (0.499 g, 5.22 mmol) in CH₃OH (15 mL). After 10 min, slowly add the resulting solution by pipette to a solution of the product of Step 1 (2.45 g, 4.97 mmol) in CH₃OH (15 mL). Monitor the reaction by TLC (15% EtOAc/toluene), and upon consumption of starting material (~1h), concentrate the mixture at room temperature in vacuo. Redissolve the resulting residue in ethyl acetate (EtOAc) and concentrate onto enough silica such that a free flowing powder is obtained. Load the resulting powder onto a chromatography column packed with 15% EtOAc/toluene. Elute with the same solvent to obtain 1.31 g (95%) of the title compound as a glass. HRMS (FAB): calcd. for M+H: C₂₆H₂₄NO₄F₂, 452.1673; found 452.1661.

25

Preparation A2

Trans-(3R,4S)-1-(4-(benzoyl)phenyl)-3-(3-phenyl)propyl]-4-(4-hydroxy)phenyl-2-azetidinone

Step 1: Reflux a mixture of 4-nitrobenzophenone (20.94 g, 92.2 mmol), ethylene glycol (25.6 mL, 461 mmol), p-toluenesulfonic acid (0.87 g, 4.61 mmol) and toluene (125 mL) overnight with azeotropic removal of water via a Dean-Stark trap. Cool the mixture to room temperature, dilute with Et₂O, wash with 1N NaOH, water and brine, dry over anhydrous Na₂SO₄ and concentrate to obtain 24.81 g (99%) of a white solid. NMR (400 MHz, CDCl₃): 8.18(2H, d, J=9.0 Hz), 7.12(2H, d, J= 9.0 Hz), 7.50(2H, d, J=8.0 Hz), 7.34(3H, m), 4.09(4H, m).

-22-

- Step 2:** Dissolve the product of step 1 (24.8 g, 92 mmol) in EtOAc (75 mL), dilute with ethanol (75 mL) and purge with N₂. Wash Raney nickel (~40 g) three times with ethanol and transfer to the reaction flask. Hydrogenate the resulting mixture on a Parr shaker at 60 psi until TLC (30% EtOAc/hexanes) indicates consumption of starting material (<2h). Filter the mixture through celite under a blanket of N₂. Wash the filter cake with 50% EtOAc/ethanol and concentrate the filtrate to give 21.6 g (97%) of a solid. NMR (400 MHz, CDCl₃): 7.50(2H, d, J=8.0 Hz), 7.30(5H, m), 6.66(2H, d, J=8.6 Hz), 4.03(4H, m).
- Step 3:** Dissolve the product of step 2 (8.49 g, 35.2 mmol) and 4-(benzyloxy)benzaldehyde (7.47 g, 35.2 mmol) in hot isopropanol (150 ml). Heat the mixture to reflux and allow the isopropanol to escape until a volume of 75 mL is obtained. Dilute the resulting solution with hexanes (200 mL) and allow to stand overnight. Collect the resultant crystals, wash with hexanes and dry under vacuum to give 14.4 g (95%) of white crystals. NMR (400 MHz, CDCl₃): 8.36(1H, s), 7.54(4H, m), 7.37(8H, m), 7.08(2H, m), 5.15(2H, s), 4.08(4H, s). MS(Cl) 436(M+H, 78), 358(39), 149(100).
- Step 4:** Add 5-phenylvaleryl chloride (10.7 mL, 53.1 mmol) to a refluxing solution of the product of step 3 (15.4 g, 35.4 mmol) and n-tributylamine (25.3 mL, 106.3 mmol) in toluene (350 mL) and reflux overnight. Cool the mixture to room temperature, quench with 1 M HCl, dilute with EtOAc, wash with 1 M HCl, NaHCO₃ (sat), water and brine, dry over anhydrous Na₂SO₄ and concentrate onto enough silica gel such that a free flowing powder results. Load the powder onto a chromatography column prepacked with 20% EtOAc/hexanes and elute with the same solvent to obtain 14 g of a solid. Recrystallize from EtOAc/hexanes to obtain 8.54 g (40%) of white crystals. NMR (400 MHz, CDCl₃): 7.30(21H, m), 6.94(2H, d, J=8.6 Hz), 5.03(2H, s), 4.54(1H, d, J=2.4 Hz), 4.01(4H, s), 3.07(1H, s), 2.63(2H, t, J=7.0 Hz), 1.92(1H, m), 1.81(3H, m).
- Step 5:** Add 6N HCl (30 mL) to a solution of the product of step 4 (4.4 g, 7.4 mmol) in THF (120 mL). After 7h, dilute with EtOAc, wash with NaHCO₃ (sat) and brine, dry over anhydrous Na₂SO₄ and concentrate to give 4.11 g (100%) of a white glass. NMR (400 MHz, CDCl₃): 7.72(4H, m), 7.55(1H, m), 7.40(8H, m), 7.27(3H, m), 7.18(3H, m), 6.98(2H, d, J=8.8 Hz), 5.05(2H, s), 4.65(1H, d, J=2.44 Hz), 3.16(1H, m),

-23-

2.65(2H, t, 7.6 Hz), 1.98(1H, m), 1.85(3H, m). HRMS(FAB) calcd for M+H, C₃₈H₃₄NO₃: 552.2539, found 552.2541.

Step 6: Add boron trichloride-dimethylsulfide (14 mL, 28.3 mmol, 2M in CH₂Cl₂) to a room temperature solution of the product of step 5 (1.56 g, 2.83 mmol) in CH₂Cl₂ (30 mL). When TLC (20% EtOAc/hexane) indicates consumption of starting material, quench the reaction by the addition of NaHCO₃ (sat). Dilute the resulting mixture with EtOAc, wash with NaHCO₃ (sat) and brine, dry over anhydrous Na₂SO₄ and concentrate onto enough silica gel such that a free flowing powder results. Load the powder onto a chromatography column prepacked with 33% EtOAc/hexanes and elute with the same solvent to obtain 1.02 g (78%) of a white glass. NMR (400 MHz, CDCl₃): 7.73(4H, m), 7.56(1H, t, 7.6 Hz), 7.45(2H, t, J=7.6 Hz), 7.34(2H, d, J=8.6 Hz), 7.28(3H, m), 7.2(2H, m), 7.16(2H, d, J=7.3 Hz), 6.85(2H, d, J=8.3 Hz), 4.65(1H, d, J=2.4 Hz), 3.15(1H, m), 2.65(2H, t, J=7.6 Hz), 1.98(1H, m), 1.85(3H, m).

Step 7: Add acetic anhydride (0.43 mL, 4.51 mmol) to a room temperature solution of the product of step 6 (1.61 g, 3.75 mmol) and N,N-dimethylaminopyridine (0.69 g, 5.64 mmol) in CH₂Cl₂ (20 mL). When TLC (30% EtOAc/hexanes) indicates consumption of starting material, dilute with EtOAc, wash with 1M HCl, water and brine, dry over anhydrous Na₂SO₄ and concentrate onto enough silica gel such that a free flowing powder results. Load the powder onto a chromatography column prepacked with 30% EtOAc/hexanes and elute with the same solvent to obtain 1.64 g (78%) of a white glass. Chiral preparative HPLC (Chiracel OD column, 20% EtOH/hexanes, 65 mL/min) provided 0.55 g of enantiomer A and 0.93 g of enantiomer B. NMR (400 MHz, CDCl₃): 7.73(4H, m), 7.56(1H, t, J=7.2 Hz), 7.46(2H, t, J=7.7 Hz), 7.32(6H, m), 7.19(3H, m), 7.12(2H, d, J=8.4 Hz), 4.70(1H, d, J=2.44 Hz), 3.17(1H, m), 2.67(2H, t, J=7.6 Hz), 2.31(3H, s), 1.97(1H, m), 1.86(3H, m). MS(CI) 504(M+H, 100), 224(100). Analytical HPLC (Chiracel OD, 20% EtOH/hexanes, 1.0 mL/min) Enantiomer A, Rt=16.83 min, Enantiomer B, Rt=23.83 min.

Step 8: Dissolve LiOH (0.098 g, 2.35 mmol) in water (2.5 mL) and add to a solution the product of step 7, enantiomer B (0.91 g, 1.8 mmol) in THF (7.5 ml). Stir overnight until TLC(30% EtOAc/hexanes) indicates consumption of starting materials. Quench the reaction with 1 M HCl,

-24-

dilute with EtOAc, wash with 1 M HCl, water and brine, dry over anhydrous Na₂SO₄ and concentrate onto enough silica gel such that a free flowing powder results. Load the powder onto a chromatography column prepacked with 30% EtOAc/hexanes and elute with the same solvent to obtain 0.36 g (46%) of a white glass. Analytical HPLC (Chiracel AS, 20% EtOH/hexanes, 0.5 mL/min), Rt=26.81 min. NMR (400 MHz, CDCl₃): 7.77(4H, m), 7.56(1H, t, J=7.6 Hz), 7.45(2H, t, J=7.6 Hz), 7.34(2H, d, J=8.6 Hz), 7.28(2H, m), 7.21(3H, m), 7.16(2H, d, J=7 Hz), 6.85(2H, d, J=8.4 Hz), 4.65(1H, d, J=2.4 Hz), 3.15(1H, m), 2.65(2H, t, J=7.4 Hz), 1.98(1H, m), 1.85(3H, m).

Preparation B

Methyl (2,3,4-tri-O-acetyl-D-glucopyransyl)uronate 1-(2,2,2-trichloroacetimidate)

Add Cs₂CO₃ (0.49 g, 1.5 mmol) to a room temperature solution of methyl 2,3,4-tri-O-acetyl-D-glucopyranuronate (5.0 g, 15 mmol) and trichloroacetonitrile (3.75 mL, 37.4 mmol) in CH₂Cl₂ (48 mL). and stir overnight. Filter the resulting brown solution through a cotton plug, washed the filtrate with water, dry over anhydrous Na₂SO₄ and concentrate. Dissolve the residue in EtOAc and concentrate onto enough silica such that a free flowing powder is obtained. Load the resulting powder onto a chromatography column packed with 30% EtOAc/hexanes. Elute with the same solvent and take only the cleanest fractions to obtain 4.35 g (61%) of the title compound as a glass. NMR (400 MHz, CDCl₃): 8.74(1H, s), 6.65(1H, d, J=3.7 Hz), 5.64(1H, t, J=9.8 Hz), 5.27(1H, t, J=9.5 Hz), 5.15(1H, dd, J=3.6, 10 Hz), 4.50(1H, d, J=10.1 Hz), 3.76(3H, s), 2.06(6H, s), 2.02(3H, s).

In a similar manner prepare:

Preparation B2

2,3,6-Tri-O-acetyl-4-O-(2,3,4,6-tetra-O-acetyl-B-D-glucopyranosyl)-α-D-glucopyranosyl 1-(2,2,2-trichloroacetimidate)

NMR (400 MHz, CDCl₃): 8.66(1H, s), 6.49(1H, d, J=3.7 Hz), 5.53(1H, t, J=10 Hz), 5.12(3H, m), 4.94(1H, t, J=8.2 Hz), 4.53(2H, m), 4.40(1H, dd, J=4.2, 12.6 Hz), 4.12(2H, m), 4.05(1H, dd, J=2.1, 12.5 Hz), 3.85(1H, t, J=9.4 Hz), 3.67(1H, m), 2.12(3H, s), 2.10(3H, s), 2.05(3H, s), 2.04(3H, s), 2.02(3H, s), 2.01(3H, s), 2.00(3H, s).

Preparation B3**2,3,4,6-Tetra-O-acetyl- α -D-glucopyranosyl 1-(2,2,2-Trichloroacetimidate)**

5 NMR (400 MHz, CDCl_3): 8.70(1H, s), 6.57(1H, d, $J=3.8$ Hz), 5.57(1H, t, $J=9.8$ Hz), 5.19(1H, t, $J=9.8$ Hz), 5.14(1H, dd, $J=3.7, 10.2$ Hz), 4.29(1H, dd, $J=4, 12.2$ Hz), 4.22(1H, m), 4.13(1H, m), 2.09(3H, s), 2.06(3H, s), 2.04(3H, s), 2.03(3H, s). MS(Electrospray): 509(M+ NH_4).

Example 1

10 **1-O-[4-[Trans-(3R,4S)-1-(4-fluorophenyl)-2-oxo-3-[3-[(S)-hydroxy-4-fluorophenyl]propyl]-4-azetidiny]phenyl]-Beta-D-glucuronic acid**

Step 1: 2,3,4-Tri-O-acetyl-1-O-[4-[trans-(3R,4S)-3-[3-[(S)-acetyloxy-3-(4-fluorophenyl)propyl-1-(4-fluorophenyl)-2-oxo-4-azetidiny]phenyl]-Beta-D-glucopyranuronic acid methyl ester

15 Add boron trifluoride etherate (0.091 mL, 0.74 mmol) to a -25 °C solution of the product of Preparation A (3.33 g, 7.38 mmol) and Preparation B (4.24 g, 8.86 mmol) in CH_2Cl_2 (74 mL) and maintain the reaction at -20 °C for 2h. Allow the reaction to warm to 10 °C over 2h. Quench the mixture with saturated NH_4Cl , dilute with EtOAc, wash with saturated NH_4Cl , water and brine, dry over anhydrous Na_2SO_4 and
20 concentrate onto enough silica such that a free flowing powder is obtained. Load the resulting powder onto a chromatography column packed with 40% EtOAc/hexanes. Elute with the same solvent to obtain 5.39 g (95%) as a foam. NMR (400 MHz, CDCl_3): 7.26(4H, m), 7.21(2H, m), 7.01 (4H, m), 6.93(2H, t, $J=8.4$ Hz), 5.69(1H, t, $J=6.7$ Hz), 5.34(2H, m), 5.29(1H, m), 5.15(1H, d, $J=7.2$ Hz), 4.56(1H, d, $J=2.1$ Hz), 4.17(1H, m), 3.73(3H, s), 3.02(1H, dt, $J=7.6, 2.3$ Hz), 2.07(14H, m), 1.85(2H, m).
25 HRMS (FAB): calcd. for M+H: $\text{C}_{39}\text{H}_{40}\text{NO}_{13}\text{F}_2$, 768.2468; found 768.2460.

30 **Step 2:** Dissolve the product of Step 1 (5.08 g, 6.98 mmol) in a mixture of CH_3OH (127 mL) and triethylamine (Et_3N) (127 mL) at room temperature. Slowly add water (445 mL) via an addition funnel over 10 min in order to maintain a homogeneous solution, then stir the resulting clear yellow solution over night. Quench a small aliquot of the reaction mixture in a vial containing 1 M HCl and EtOAc and monitor
35 consumption of the starting material by TLC (5% acetic acid (HOAc)/20% CH_3OH /75% CH_2Cl_2) of the EtOAc layer. Remove the CH_3OH and Et_3N

-26-

on a rotary evaporator, acidify the remaining solution with 1M HCl, dilute with EtOAc and extract with EtOAc. Combine the extracts, wash with 1M HCl, water and brine, dry over anhydrous Na₂SO₄ and concentrate to a white solid 3.82 g (93%). Dissolve the solid in CH₂Cl₂, and concentrate onto enough silica such that a free flowing powder is obtained. Load the resulting powder onto a chromatography column packed with silica and 15% CH₃OH/CH₂Cl₂. Elute with 5% HOAc/15% CH₃OH/80% CH₂Cl₂. Concentrate the fractions containing the title compound, azeotrope first with toluene (3X) and then CH₃OH (5X). Heat the resultant solid to 60°C under vacuum to remove any residual solvent and obtain the title compound as a white solid 2.6 g (64%). NMR (400 MHz, CD₃OD): 7.29(6H, m), 7.09(1H, d, J=8.6 Hz), 6.70(4H, m), 4.96(1H, m), 4.80(1H, d, J=2.0 Hz), 4.59(1H, m), 3.97(1H, d, J=9.6 Hz), 3.59(1H, m), 3.49(2H, m), 3.09(1H, m), 1.86(4H, m). HRMS (FAB): calcd. for M+H: C₃₀H₃₀NO₉F₂, 586.1889; found 586.1883.

Example 1A

1-O-[4-[Trans-(3R,4S)-1-(4-iodophenyl)-2-oxo-3-[3-[(S)-hydroxy-4-fluorophenyl]propyl]-4-azetidiny]phenyl]-Beta-D-glucuronic acid

Treat 1-(4-iodophenyl)-3(R)-[3(S)-acetyloxy-3-(4-fluorophenyl)-propyl]-4(S)-(4-hydroxyoxyphenyl)-2-azetidinone and the product of Preparation B according to the procedure described in Example 1 to obtain the title compound. M.p. 135-137 °C; FAB MS calc'd for C₃₀H₂₉FINO₉ NaCl m/z = 751.05, found m/z = 751.2.

Example 2

1-O-[4-[Trans-(3R,4S)-1-(4-fluorophenyl)-2-oxo-3-[3-[(S)-hydroxy-4-fluorophenyl]propyl]-4-azetidiny]phenyl]-3-O-(Beta-D-glucopyranosyl)-Beta-D-glucopyranose

Step 1: 2,3,6-Tri-O-acetyl-4-O-(2,3,4,6-tetra-O-acetyl-B-D-glucopyranosyl)-1-O-[4-[trans-(3R,4S)-3-[3(S)-acetyloxy-3-(4-fluorophenyl)-propyl]-1-(4-fluorophenyl)-2-oxo-4-azetidiny]phenyl]-Beta-D-glucopyran

Using a procedure similar to that described in Example 1, Step 1, combine the product of Preparation A and Preparation B2 to obtain the title compound of Step 1. NMR (400 MHz, CDCl₃): 7.23(6H, m), 6.97(6H, m), 5.69(1H, t, 6.6 Hz), 5.26(1H, t, J=9.1 Hz), 5.11(4H, m), 4.95(1H, t, J=8.2 Hz), 4.54(3H, m), 4.39(1H, dd, J=4.3, 12.5 Hz), 4.06(2H, m), 3.87(1H, t, J=9.5 Hz), 3.75(1H, m), 3.68(1H, m), 3.02(1H, dt, J=2.1,

-27-

7.6 Hz), 2.05(26H, m), 1.85(2H, m). HRMS (FAB): calcd. for M+Na: C₅₂H₅₇NO₂₁F₂Na, 1092.3289; found 1092.3308.

Step 2: Using a procedure similar to that described in Example 1, Step 2, treat the product of Step 1, above, to obtain the title compound of Example 2. NMR (400 MHz, CD₃OD): 7.29(6H, m), 7.10(2H, d, J=8.7 Hz), 7.01(4H, m), 4.96(1H, under CD₃OD), 4.81(1H, d, J=2.2 Hz), 4.60(1H, m), 4.43(1H, d, J=7.9 Hz), 3.88(3H, m), 3.62(4H, m), 3.51(1H, d, J=8.9 Hz), 3.34(2H, m), 3.24(1H, t, J=8.8 Hz), 3.08(1H, m), 1.88(7H, m). MS (FAB): 756 (M+Na, 70), 734(M+, 100), 716(716, 20).

10

Example 3

1-O-[4-[Trans-(3R,4S)-3-[3(S)-hydroxy-3-(4-fluorophenyl)propyl-1-(4-fluorophenyl)-2-oxo-4-azetidiny]]phenyl]-Beta-D-glucopyranose

Step 1: 2,3,4,5-Tetra-O-acetyl-1-O-[4-[trans-(3R,4S)-3-[3(S)-acetyloxy-3-(4-fluorophenyl)propyl-1-(4-fluorophenyl)-2-oxo-4-azetidiny]]phenyl]-Beta-D-glucopyran

Using a procedure similar to that described in Example 1, Step 1, combine the product of Preparation A and Preparation B3 to obtain the title compound of Step 1. NMR (400 MHz, CDCl₃): 7.26(4H, m), 7.20(2H, m), 7.01(4H, m), 6.93(2H, t, J=8.5 Hz), 5.69(1H, t, J=6.5 Hz), 5.29(2H, m), 5.18(1H, t, J=9.7 Hz), 5.09(1H, d, J=7.3 Hz), 4.56(1H, d, J=2.2 Hz), 4.29(1H, dd, J=5.2, 12.2 Hz), 4.17(1H, dd, J=2.2 Hz, 12.2 Hz), 3.85(1H, m), 3.03(1H, dt, J=2.1, 7.5 Hz), 2.06(17H, m), 1.85 (2H, m). HRMS (FAB): calcd. for M+Na: C₄₀H₄₁NO₁₃F₂Na, 804.2444, found 804.2432.

Step 2: Using a procedure similar to that described in Example 1, Step 2, treat the product of Step 1, above, to obtain the title compound of Example 3. NMR (400 MHz, CD₃OD): 7.29(6H, m), 7.11(2H, d, J=8.8 Hz), 6.98(4H, m), 4.89(1H, under CD₃OD), 4.80(1H, d, J=2.2 Hz), 4.60(1H, m), 3.88(1H, dd, J=2.0, 12.0 Hz), 3.68(1H, dd, J=5.4, 12.0 Hz), 3.41(3H, m), 3.08(1H, m), 1.86(4H, m). MS (FAB): 572 (M+H, 40), 392(100).

30

Example 4

1-O-[4-[Trans-(3R,4S)-1-(4-fluorophenyl)-2-oxo-3-[3-(S)-hydroxy-4-fluorophenyl)propyl]]-4-azetidiny]]phenyl]-Beta-D-glucuronic acid methyl ester

Add KCN (0.028 g, 0.43 mmol) to a room temperature solution of the product of Example 1, Step 1, (0.312 g, 0.43 mmol) in CH₃OH (5 mL) and stir the mixture overnight. Monitor by TLC (10% CH₃OH/CH₂Cl₂);

35

-28-

heat the mixture to 40 °C for 2.5 h. Cool the mixture to room temperature, concentrate onto enough silica such that a free flowing powder is obtained. Load the resulting powder onto a chromatography column packed with silica and 5% CH₃OH/CH₂Cl₂. Elute with 5%
5 CH₃OH/CH₂Cl₂ and collect the purest fractions to obtain 0.116 g of the title compound. NMR (400 MHz, CDCl₃/CD₃OD): 7.16(6H, m), 6.95(4H, m), 6.86(2H, t, J=8.6 Hz), 4.83(1H, d, J=7.6 Hz), 4.56(1H, t, J=6.3 Hz), 4.55(1H, d, J=2.1 Hz), 3.90(1H, d, J=9.8 Hz), 3.73(3H, s), 3.67(1H, t, J=9.1 Hz), 3.51(1H, m), 3.46(1H, t, J=9.2 Hz), 3.30(1H, s), 2.98(1H, m),
10 1.80(4H, m). HRMS (FAB): calcd. for M+H: C₃₁H₃₂NO₉F₂, 600.2045; found 600.2049.

Example 5

1-O-[4-[Trans-(3R,4S)-1-(4-methoxyphenyl)-2-oxo-3-(3-phenyl)propyl]-4-azetidiny]phenyl]-Beta-D-glucuronic acid methyl ester

15 Step 1: 2,3,4-Tri-O-acetyl-1-O-[4-[Trans-(3R,4S)-3-[3-[(S)-acetyloxy-3-(4-fluorophenyl)propyl]-1-(4-methoxyphenyl)-2-oxo-4-azetidiny]phenyl]-Beta-D-glucopyranuronic Acid Methyl Ester

Add triphenylphosphine (0.19 g, 0.72 mmol) to a 0 °C solution of 1,1'-(azodicarbonyl)dipiperidine (0.18 g, 0.72 mmol) in THF (3 mL). After
20 10 min, add (3R,4S)-4-(4-hydroxyphenyl)-1-(4-methoxyphenyl)-3-(3-phenylpropyl)-2-azetidinone (0.2 g, 0.52 mmol), followed by methyl-2,3,4-tri-O-acetyl-D-glucopyranuronate (0.21 g, 0.62 mmol). Allow the mixture to warm to room temperature overnight. Concentrate the mixture onto enough silica such that a free flowing powder is obtained. Load the
25 resulting powder onto a chromatography column packed with silica and 30% EtOAc/hexanes. Elute with 30-50% EtOAc/hexanes to obtain 0.198 g of material which is further purified by silica chromatography eluting with 20% CH₃OH/CH₂Cl₂ to provide 0.074 g of the title compound of Step 1. NMR (400 MHz, CDCl₃): 7.27(4H, m), 7.17(5H, m), 6.98(2H, J=8.5 Hz), 6.77(2H, m), 5.30(3H, m), 5.13(1H, d, J=7.3 Hz), 4.56(1H, d, J=1.9 Hz), 4.17(1H, m), 3.74(3H, s), 3.73(3H, s), 3.04(1H, m), 2.64(2H, t, J=7.6 Hz), 2.05(9H, m), 1.97(1H, m), 1.82(3H, m). HRMS (FAB): calcd. for M+H: C₃₈H₄₂NO₁₂ 704.2707; found 704.2696.

Step 2: Using a procedure similar to that of Example 4, treat the product
35 of Step 1 to obtain the title compound. NMR (400 MHz, CDCl₃): 7.27(4H, m), 7.17(5H, m), 7.04(2H, J=8.6 Hz), 6.75(2H, J=9.1 Hz),

4.90(1H, d, J=7.0 Hz), 4.55(1H, d, J=1.8 Hz), 3.98(1H, d, J=9.7 Hz), 3.88(1H, t, J=8.6 Hz), 3.76(8H, m), 3.03(1H, m), 2.63(2H, t, J=6.7 Hz), 1.95(1H, m), 1.81(3H, m). HRMS (FAB): calcd. for M+H: C₃₂H₃₆NO₉, 578.2390; found 578.2379.

5

Example 6

1-O-[4-[Trans-(3R,4S)-1-(4-(benzoyl)phenyl)-2-oxo-3-(3-phenyl)propyl]-4-azetidiny]phenyl]-Beta-D-glucuronic acid methyl ester

Step 1: 2,3,4-Tri-O-acetyl-1-O-[4-[Trans-(3R,4S)-1-(4-(benzoyl)phenyl)-2-oxo-3-(3-phenyl)propyl]-4-azetidiny]phenyl]-Beta-D-glucuronic acid methyl ester

10

In a similar manner to Example 5, Step 1, treat (3R,4S)-1-(4-benzoylphenyl)-4-(4-hydroxyphenyl)-3-(3-phenylpropyl)-2-azetidinone and methyl-2,3,4-tri-O-acetyl-D-glucopyranuronate to obtain the title compound of Step 1. NMR (400 MHz, CDCl₃): 7.73(4H, m), 7.57(1H, t, J=7.0 Hz), 7.46(2H, t, J=8.0 Hz), 7.30(6H, m), 7.21(1H, d, J=7.1 Hz), 7.16(2H, d, J=8.0 Hz), 7.01(2H, d, J=8.5 Hz), 5.31(3H, m), 5.15(1H, d, J=7.3 Hz), 4.67(1H, d, J=2.2 Hz), 4.17(1H, dd, J=2.7, 6.7 Hz), 3.73(3H, s), 3.14(1H, m), 2.66(2H, t, J=7.4 Hz), 2.06(9H, m), 1.98(1H, m), 1.85(3H, m). HRMS (FAB): calcd. for M+H: C₄₄H₄₄NO₁₂, 778.2864; found 778.2849.

15

Step 2: Using a procedure similar to that of Example 4, treat the product of Step 1 to obtain the title compound. NMR (400 MHz, CDCl₃): 7.72(2H, overlapping d, J=8.6, 7.6 Hz), 7.56(1H, t, J=7.6 Hz), 7.45(2H, t, J=7.7 Hz), 7.30(6H, m), 7.20(1H, d, J=7.0 Hz), 7.16(2H, d, J=7.6 Hz), 7.08(2H, d, J=8.6 Hz), 4.93(1H, d, J=7.0 Hz), 4.67(1H, dd, J=2.1 Hz), 3.99(1H, d, J=9.8 Hz), 3.88(1H, t, J=8.6 Hz), 3.81(3H, s), 3.73(2H, m), 3.14(1H, m), 2.65(2H, t, J=7.6 Hz), 1.98(1H, m), 1.84(3H, m). HRMS (FAB): calcd. for M+H: C₃₈H₃₈NO₉, 652.2547; found 652.2528.

25

Example 7

1-O-[4-[Trans-(3R,4S)-1-(4-methoxyphenyl)-2-oxo-3-(3-phenylpropyl)-4-azetidiny]phenyl]-Beta-D-glucopyranose

30

Step 1: 1-O-[4-[Trans-(3R,4S)-1-(4-methoxyphenyl)-2-oxo-3-(3-phenylpropyl)-4-azetidiny]phenyl]-2,3,4,6,-tetra-O-(phenylmethyl)-Beta-D-glucopyranose

35

Add n-tributylphosphine (1.45 mL, 5.81 mmol) to a 0 °C solution of 1,1'-(azodicarbonyl)dipiperidine (1.47 g, 5.81 mmol) in THF (30 mL). After 5 min., add (3R,4S)-4-(4-hydroxyphenyl)-1-(4-methoxy-phenyl)-3-

-30-

- (3-phenylpropyl)-2-azetidinone (1.5 g, 3.87 mmol), followed by 2,3,4,6-tetra-O-benzyl-D-glucopyranose (2.72 g, 5.03 mmol). The reaction becomes very thick, and additional THF (30 mL) is added to facilitate stirring; the mixture is allowed to warm to room temperature overnight.
- 5 Filter the mixture through celite, wash the filter cake with EtOAc, and concentrate the filtrate onto enough silica such that a free flowing powder is obtained. Load the resulting powder onto a chromatography column packed with silica and 5% EtOAc/toluene. Elute with the same solvent to obtain 3.57 g (~100%) of the title compound of Step 1 as a
- 10 thick syrup. NMR (400 MHz, CDCl₃): 7.16(19H, m), 7.19(10H, m), 7.04(2H, d, J=8.7 Hz), 6.76(2H, d, J=9.2 Hz), 4.98(3H, m), 4.83(3H, m), 4.55(4H, m), 3.70(9H, m), 3.05(1H, m), 2.65(2H, t, J=7.3 Hz), 1.96(1H, m), 1.83(3H, m). MS (FAB): 910(M⁺, 55), 568(40), 478(100), 386(55).
- 15 Step 2: Dissolve the product of Step 1 (0.20 g, 0.35 mmol) in CH₃OH (4.5 mL), dilute with EtOAc (4.5 mL) and purge with nitrogen. Add 20% Pd(OH)₂ on carbon (0.35 g), purge the resulting mixture with hydrogen (3X) and then stir under a balloon of hydrogen overnight. Filter the mixture through celite and wash the filter cake with EtOAc followed by CH₃OH. Concentrate the filtrate to a clear foam 0.161 g (83% crude).
- 20 Purify the foam further by silica chromatography eluting with 5% CH₃OH/EtOAc to obtain 0.127 g (66%) of the the title compound as a white powder. NMR (400 MHz, CD₃OD): 7.18(11H, m), 6.78(2H, d, J=8.9 Hz), 4.88(1H, partially obscured by CD₃OD), 4.72(1H, d, J=1.2 Hz) 3.88(1H, d, J=11.7 Hz), 3.70(4H, m), 3.41(4H, m), 3.03(1H, m), 2.60(2H,
- 25 t, J=7.0 Hz), 1.79(4H, m). HRMS (FAB): calcd. for M+H: C₃₁H₃₆NO₈, 550.2441; found 500.2424.

Example 8

1-O-[4-[Trans-(3R,4S)-1-(4-methoxyphenyl)-2-oxo-3-(3-phenylpropyl)-4-azetidiny]]phenyl]-Beta-D-glucuronic acid

- 30 Step 1: 2,3,4-tri-O-Benzyl-1-O-[4-[trans-(3R,4S)-1-(4-fluorophenyl)-2-oxo-3-[3-[(S)-hydroxy-4-fluorophenyl]propyl]]-4-azetidiny]]phenyl]-Beta-D-glucuronic acid benzyl ester

- 35 Use (3R,4S)-4-(4-hydroxyphenyl)-1-(4-methoxy-phenyl)-3-(3-phenylpropyl)-2-azetidinone and benzyl 2,3,4-tri-O-benzyl-D-glucopyranuronate in a procedure similar to that described in Example 7, Step 1, to obtain the title compound of Step 1. NMR (400 MHz, CDCl₃):

7.22(29H, m), 7.01(2H, d, J=8.7 Hz), 6.77(2H, d, J=9.1 Hz), 5.15(2H, app. d, J=3.8 Hz), 5.01(1H, d, J=7.2 Hz), 4.97(1H, d, J=11 Hz), 4.90(1H, d, J=11 Hz), 4.80(2H, d, J=11 Hz), 4.74(1H, d, J=10.7 Hz), 4.56(1H, d, J=2.2 Hz), 4.50(1H, d, J=10.7 Hz), 4.04(1H, d, J=9.6 Hz), 3.93(1H, t, J=8.6 Hz), 3.73(5H, m), 3.05(1H, m), 2.65(2H, t, J=7.6 Hz), 1.96(1H, m), 1.83(3H, m). HRMS (FAB): calcd. for M+H: C₅₉H₅₈NO₉ 924.4112; found 924.4119.

Step 2: Using a procedure similar to Example 7, Step 2, treat the product of Step 1 to obtain the title compound of Example 8. NMR (400 MHz, CD₃OD): 7.31(2H, d, J=8.9 Hz), 7.21(7H, m), 7.09(2H, d, J=8.7 Hz), 7.81(2H, d, J=8.9 Hz), 4.97(1H, dd, J=1.9, 5.5 Hz), 4.76(1H, d, J=2.0 Hz), 3.97(1H, d, J=9.7 Hz), 3.72(3H, s), 3.60(1H, m), 3.49(2H, m), 3.08(1H, m), 2.64(2H, t, J=7.2 Hz), 1.83(4H, m). HRMS (FAB): calcd. for M+H: C₃₁H₃₄NO₉ 564.2234; found 564.2242.

15

Example 9

1-Methyl-6-O-[4-[trans-(3R,4S)-1-(4-methoxyphenyl)-2-oxo-3-(3-phenylpropyl)-4-azetidiny]phenyl]-Alpha-D-glucopyranoside

Step 1: 1-Methyl-2,3,4-tri-O-Benzyl-6-O-[4-[Trans-(3R,4S)-1-(4-methoxyphenyl)-2-oxo-3-(3-phenylpropyl)-4-azetidiny]phenyl]-Alpha-D-glucopyranoside

20

Use (3R,4S)-4-(4-hydroxyphenyl)-1-(4-methoxy-phenyl)-3-(3-phenylpropyl)-2-azetidinone and methyl 2,3,4-tri-O-benzyl-D-glucopyranoside in a procedure similar to that described in Example 7, Step 1, to obtain the title compound of Step 1. NMR (400 MHz, CDCl₃): 7.26(24H, m), 6.85(2H, d, J=8.6 Hz), 6.74(2H, d, J=9 Hz), 5.01(1H, d, J=10.7 Hz), 4.86(1H, d, J=11.0 Hz), 4.85(1H, d, J=10.7 Hz), 4.82(1H, d, J=12.1 Hz), 4.69(1H, d, J=12.1 Hz), 4.63(1H, d, J=3.6 Hz), 4.54(1H, d, J=2.3 Hz), 4.51(1H, d, J=11.0 Hz), 4.09(2H, d, J=2.8 Hz), 4.03(1H, t, J=9.6 Hz), 3.90(1H, d, J=10.1 Hz), 3.72(3H, s), 3.60(1H, dd, J=3.6, 9.6 Hz), 3.38(3H, s), 3.06(1H, m), 2.64(2H, t, J=7.6 Hz), 1.97(1H, m), 1.83(3H, m).

30

Step 2: Using a procedure similar to Example 7, Step 2, treat the product of Step 1 to obtain the title compound of Example 9. NMR (400 MHz, CDCl₃): 7.22(9H, m), 6.94(2H, d, J=8.6 Hz), 6.76(2H, d, J=8.9 Hz), 4.81(1H, d, J=3.9 Hz), 4.54(1H, d, J=2.2 Hz), 4.22(2H, m), 3.97(1H, m), 3.71(5H, m), 3.56(1H, dd, J=3.9, 9.1 Hz), 3.44(3H, s), 3.06(1H, m),

35

-32-

2.64(2H, d, J=7.4 Hz), 1.91(1H, m), 1.82(3H, m). HRMS (FAB): calcd. for M+H: C₃₂H₃₈NO₈ 564.2597, found 564.2578.

Example 10

5 1-O-[4-[Trans-(3R,4S)-1-(4-(benzoyl)phenyl)-2-oxo-3-(3-phenyl)propyl]-4-azetidiny]phenyl]-Beta-D-glucuronic acid

Add LiOH (0.6 mL, 0.6 mmol, 1M) to a room temperature solution of the product of Example 6 (0.064 g, 0.1 mmol) in THF (2 mL). After 50 min., dilute the mixture with EtOAc, quench with HCl (1M), wash with HCl (1M) and brine, dry over anhydrous Na₂SO₄ and concentrate to a white foam 0.60 g (97%). NMR (400 MHz, CD₃OD): 7.67(4H, m), 7.60(1H, m), 7.48(3H, m), 7.36(2H, d, J=8.8 Hz), 7.34(2H, d, J=8.8 Hz), 7.23(2H, m), 7.14(2H, d, J=7.5 Hz), 7.10(2H, d, J=8.7 Hz), 4.97(1H, m), 4.87(1H, d, J=2.2 Hz), 3.97(1H, d, J=9.7 Hz), 3.60(1H, m), 3.49(2H, m), 3.17(1H, m), 2.63(2H, t, J=7.4 Hz), 1.89(1H, m), 1.81(3H, m). HRMS (FAB): calcd. for M+H: C₃₇H₃₆NO₉ 638.2390; found 638.2377.

Example 11

1-O-[4-[Trans-(3R,4S)-1-(4-fluorophenyl)-2-oxo-3-[3-[(S)-hydroxy-4-iodophenyl]propyl]-4-azetidiny]phenyl]-Beta-D-glucuronic acid
Step 1: Condense 1-(4-fluorophenyl)-3(R) -[3(S)-acetyloxy-3-(4-bromophenyl)propyl]-4(S)-(4-hydroxyphenyl)-2-azetidinone and the product of Preparation B with boron trifluoride etherate according to the procedure described in Example 1. To a solution of the resultant tetraacetate (250 mg, 0.30 mmol) in CH₃OH (2 mL) cooled to 0°C, add KCN (10 mg, 0.15 mmol) and stir at room temperature for 2h, then heat to 45°C for 4.5h. Cool the mixture to room temperature and partition between water (20mL) and EtOAc (30 mL). Wash the EtOAc layer with water and brine, dry (Na₂SO₄) and concentrate in vacuo. Adsorb the residue (230 mg) onto SiO₂ and chromatograph over SiO₂ (25 g), eluting with 2% CH₃OH in CH₂Cl₂ progressing to 10% CH₃OH in CH₂Cl₂ to give, after concentration, 84 mg (43%) of the aryl bromide as a solid.

Step 2: To the product of step 1 (25 mg, 0.038 mmol) dissolved in degassed DMF (0.4 mL), add hexabutylditin (220 mg, 38 mmol) and tetrakis triphenylphosphine palladium (4.4 mg, 0.0038 mmol) and heat the mixture to 95°C under argon for 5h. Cool the reaction, concentrate in vacuo and adsorb the resulting residue directly onto SiO₂.

-33-

Chromatograph over SiO₂ (4g), eluting with CH₂Cl₂ progressing to 10% CH₃OH in CH₂Cl₂. Rechromatograph the desired fraction as above and after concentration obtain 7.4 mg (22%) of the desired aryl stannane as a waxy solid.

- 5 **Step 3:** To the product of step 2 (11.8 mg, 0.0135 mmol) dissolved in CH₃OH (2 mL) containing pH 5.8 phosphate buffer (0.3mL), add a 1M solution of NaI in water (14mL, 0.014 mmol). To this mixture add 68 iodobeads® (~37 mmol) and gently shake the resulting mixture for 1.5h at room temperature. Filter the iodobeads and wash with EtOH and a
10 small amount of ether. Concentrate the filtrate and partition the residue between EtOAc and 10% aqueous Na₂SO₃, dry the EtOAc layer (MgSO₄) and concentrate in vacuo. Adsorb the residue onto SiO₂ and chromatograph over SiO₂ (2g), eluting with CH₂Cl₂ progressing to 6% CH₃OH in CH₂Cl₂. Concentrate the appropriate fractions to obtain 6.1
15 mg (64%) of the methyl ester of the title compound as a solid.

- Step 4:** Stir a solution of the product of step 3 (6.1 mg, 8.6mmol) in a mixture of water (0.7mL), triethylamine (0.2 mL) and CH₃OH (0.1 mL) at room temperature for 30 min. Concentrate the mixture in vacuo to give 5 mg (83%) of the title compound as a solid. M.p. 157 - 159 °C, FAB MS
20 calc'd for C₃₀H₃₀FINO₉ m/z = 694.1, found m/z = 694.1.

The following formulations exemplify some of the dosage forms of this invention. In each, the term "active compound" designates a compound of formula I.

EXAMPLE A - Tablets

<u>No.</u>	<u>Ingredient</u>	<u>mg/tablet</u>	<u>mg/tablet</u>
1	Active Compound	100	500
2	Lactose USP	122	113
3	Corn Starch, Food Grade, as a 10% paste in Purified Water	30	40
4	Corn Starch, Food Grade	45	40
5	Magnesium Stearate	3	7
Total		300	700

25 Method of Manufacture

Mix Item Nos. 1 and 2 in suitable mixer for 10-15 minutes.

Granulate the mixture with Item No. 3. Mill the damp granules through a coarse screen (e.g., 1/4", 0.63 cm) if necessary. Dry the damp granules.

-34-

Screen the dried granules if necessary and mix with Item No. 4 and mix for 10-15 minutes. Add Item No. 5 and mix for 1-3 minutes. Compress the mixture to appropriate size and weight on a suitable tablet machine.

EXAMPLE B - Capsules

<u>No.</u>	<u>Ingredient</u>	<u>mg/tablet</u>	<u>mg/tablet</u>
1	Active Compound	100	500
2	Lactose USP	106	123
3	Corn Starch, Food Grade	40	70
4	Magnesium Stearate NF	4	7
	Total	250	700

5 Method of Manufacture

Mix Item Nos. 1, 2 and 3 in a suitable blender for 10-15 minutes. Add Item No. 4 and mix for 1-3 minutes. Fill the mixture into suitable two-piece hard gelatin capsules on a suitable encapsulating machine.

Representative formulations comprising a cholesterol biosynthesis inhibitor are well known in the art. It is contemplated that where the two active ingredients are administered as a single composition, the dosage forms disclosed above for substituted azetidinone compounds may readily be modified using the knowledge of one skilled in the art.

15 The in vivo activity of the compounds of formula I can be determined by the following procedure.

In Vivo Assay of Hypolipidemic Agents Using the Hyperlipidemic Hamster

Hamsters are separated into groups of six and given a controlled cholesterol diet (Purina Chow #5001 containing 0.5% cholesterol) for seven days. Diet consumption is monitored to determine dietary cholesterol exposure in the presence of test compounds. The animals are dosed with the test compound once daily beginning with the initiation of diet. Dosing is by oral gavage of 0.2mL of corn oil alone (control group) or solution (or suspension) of test compound in corn oil. All animals moribund or in poor physical condition are euthanized. After seven days, the animals are anesthetized by IM injection of ketamine and sacrificed by decapitation. Blood is collected into Vacutainer™ tubes containing EDTA for plasma total cholesterol and triglyceride analysis and the liver excised for free and esterified

cholesterol and triglyceride tissue analysis. Data is reported as percent reduction of plasma cholesterol and hepatic cholesterol esters versus control levels.

Using the test procedures described above, the following

5 in vivo data were obtained for compounds of formula I. Data is reported as percent change (i.e., percent reduction in plasma cholesterol and in hepatic cholesterol esters) versus control, therefore, negative numbers indicate a positive cholesterol-lowering effect. For racemic compounds of formula I or active diastereomers or enantiomers of compounds of


10 formula I, compounds administered at dosages of 3 to 10 mg/kg show a range of 0 to -98% reduction in hepatic cholesterol esters, while compounds administered at dosages of 0.01 to 1 mg/kg show a range of -19 to -94% reduction in hepatic cholesterol esters. Compounds preferably show a range of -50 to -98% reduction in hepatic cholesterol

15 esters at a dosage range of 0.01 to 1 mg/kg.

-36-

We claim:



1. A compound represented by the structural formula



- 5 or a pharmaceutically acceptable salt thereof, wherein

R^{26} is H or OG^1 ;

G and G^1 are independently selected from the group consisting of



provided that when R^{26} is H or

OH, G is not H;

- 10 R, R^a and R^b are independently selected from the group consisting of H, -OH, halogeno, -NH₂, azido, (C_1-C_6) alkoxy (C_1-C_6) -alkoxy or -W- R^{30} ;

W is independently selected from the group consisting of -NH-C(O)-, -O-C(O)-, -O-C(O)-N(R^{31})-, -NH-C(O)-N(R^{31})- and -O-C(S)-N(R^{31})-;

- 15 R^2 and R^6 are independently selected from the group consisting of H, (C_1-C_6) alkyl, aryl and aryl (C_1-C_6) alkyl;

R^3 , R^4 , R^5 , R^7 , R^{3a} and R^{4a} are independently selected from the group consisting of H, (C_1-C_6) alkyl, aryl (C_1-C_6) alkyl, -C(O) (C_1-C_6) alkyl and -C(O)aryl;

- 20 R^{30} is independently selected from the group consisting of R^{32} -substituted T, R^{32} -substituted-T- (C_1-C_6) alkyl, R^{32} -substituted- $(C_2-$

-37-

C₄)alkenyl, R³²-substituted-(C₁-C₆)alkyl, R³²-substituted-(C₃-C₇)cycloalkyl and R³²-substituted-(C₃-C₇)cycloalkyl(C₁-C₆)alkyl;

R³¹ is independently selected from the group consisting of H and (C₁-C₄)alkyl;


5 T is independently selected from the group consisting of phenyl, furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl and pyridyl;

R³² is independently selected from 1-3 substituents independently selected from the group consisting of halogeno, (C₁-C₄)alkyl, -OH, phenoxy, -CF₃, -NO₂, (C₁-C₄)alkoxy, methylenedioxy, 10 oxo, (C₁-C₄)alkylsulfanyl, (C₁-C₄)alkylsulfinyl, (C₁-C₄)alkylsulfonyl, -N(CH₃)₂, -C(O)-NH(C₁-C₄)alkyl, -C(O)-N((C₁-C₄)alkyl)₂, -C(O)-(C₁-C₄)alkyl, -C(O)-(C₁-C₄)alkoxy and pyrrolidinylcarbonyl; or R³² is a covalent bond and R³¹, the nitrogen to which it is attached and R³² form 15 a pyrrolidinyl, piperidinyl, N-methyl-piperazinyl, indolinyl or morpholinyl group, or a (C₁-C₄)alkoxycarbonyl-substituted pyrrolidinyl, piperidinyl, N-methylpiperazinyl, indolinyl or morpholinyl group;

Ar¹ is aryl or R¹⁰-substituted aryl;

Ar² is aryl or R¹¹-substituted aryl;

20 Q is a bond or, with the 3-position ring carbon of the azetidinone,



R¹ is selected from the group consisting of


-(CH₂)_q-, wherein q is 2-6, provided that when Q forms a spiro ring, q can also be zero or 1;

25 -(CH₂)_e-E-(CH₂)_r-, wherein E is -O-, -C(O)-, phenylene, -NR²²- or -S(O)₀₋₂-, e is 0-5 and r is 0-5, provided that the sum of e and r is 1-6;

-(C₂-C₆)alkenylene-; and

30 -(CH₂)_f-V-(CH₂)_g-, wherein V is C₃-C₆ cycloalkylene, f is 1-5 and g is 0-5, provided that the sum of f and g is 1-6;

R¹² is




-38-

R¹³ and R¹⁴ are independently selected from the group consisting of -CH₂-, -CH(C₁-C₆ alkyl)-, -C(di-(C₁-C₆) alkyl)-, -CH=CH- and -C(C₁-C₆ alkyl)=CH-; or R¹² together with an adjacent R¹³, or R¹² together with an adjacent R¹⁴, form a -CH=CH- or a -CH=C(C₁-C₆ alkyl)- group;

a and b are independently 0, 1, 2 or 3, provided both are not zero; provided that when R¹³ is -CH=CH- or -C(C₁-C₆ alkyl)=CH-, a is 1; provided that when R¹⁴ is -CH=CH- or -C(C₁-C₆ alkyl)=CH-, b is 1; provided that when a is 2 or 3, the R¹³'s can be the same or different; and provided that when b is 2 or 3, the R¹⁴'s can be the same or different;

and when Q is a bond, R¹ also can be:



M is -O-, -S-, -S(O)- or -S(O)₂-;

X, Y and Z are independently selected from the group consisting of -CH₂-, -CH(C₁-C₆)alkyl- and -C(di-(C₁-C₆)alkyl);

R¹⁰ and R¹¹ are independently selected from the group consisting of 1-3 substituents independently selected from the group consisting of (C₁-C₆)alkyl, -OR¹⁹, -O(CO)R¹⁹, -O(CO)OR²¹, -O(CH₂)₁₋₅OR¹⁹, -O(CO)NR¹⁹R²⁰, -NR¹⁹R²⁰, -NR¹⁹(CO)R²⁰, -NR¹⁹(CO)OR²¹, -NR¹⁹(CO)NR²⁰R²⁵, -NR¹⁹SO₂R²¹, -COOR¹⁹, -CONR¹⁹R²⁰, -COR¹⁹, -SO₂NR¹⁹R²⁰, S(O)₀₋₂R²¹, -O(CH₂)₁₋₁₀-COOR¹⁹, -O(CH₂)₁₋₁₀CONR¹⁹R²⁰, -(C₁-C₆ alkylene)-COOR¹⁹, -CH=CH-COOR¹⁹, -CF₃, -CN, -NO₂ and halogen;

R¹⁵ and R¹⁷ are independently selected from the group consisting of -OR¹⁹, -O(CO)R¹⁹, -O(CO)OR²¹ and -O(CO)NR¹⁹R²⁰; R¹⁶ and R¹⁸ are independently selected from the group consisting of H, (C₁-C₆)alkyl and aryl; or R¹⁵ and R¹⁶ together are =O, or R¹⁷ and R¹⁸ together are =O;

d is 1, 2 or 3;

h is 0, 1, 2, 3 or 4;


s is 0 or 1; t is 0 or 1; m, n and p are independently 0-4; provided that at least one of s and t is 1, and the sum of m, n, p, s and t is 1-6;

-39-

provided that when p is 0 and t is 1, the sum of m, s and n is 1-5; and
provided that when p is 0 and s is 1, the sum of m, t and n is 1-5;

v is 0 or 1;

j and k are independently 1-5, provided that the sum of j, k and v
5 is 1-5;



and when Q is a bond and R¹ is , Ar¹ can
also be pyridyl, isoxazolyl, furanyl, pyrrolyl, thienyl, imidazolyl, pyrazolyl,
thiazolyl, pyrazinyl, pyrimidinyl or pyridazinyl;

R¹⁹ and R²⁰ are independently selected from the group
10 consisting of H, (C₁-C₆)alkyl, aryl and aryl-substituted (C₁-C₆)alkyl;
R²¹ is (C₁-C₆)alkyl, aryl or R²⁴-substituted aryl;
R²² is H, (C₁-C₆)alkyl, aryl (C₁-C₆)alkyl, -C(O)R¹⁹ or -COOR¹⁹;
R²³ and R²⁴ are independently 1-3 groups independently
selected from the group consisting of H, (C₁-C₆)alkyl, (C₁-C₆)alkoxy,
15 -COOH, NO₂, -NR¹⁹R²⁰, -OH and halogeno; and
R²⁵ is H, -OH or (C₁-C₆)alkoxy.

2. A compound of claim 1 wherein:

Ar¹ is phenyl or halogeno-substituted phenyl;

20 Ar² is phenyl, lower alkoxy-substituted phenyl or halogeno-
substituted phenyl;

Q is a bond and R¹ is lower alkylene;

Q, with the 3-position ring carbon of the azetidinone, forms the

group  wherein R¹³ and R¹⁴ are each ethylene and a

25 and b are each 1, and wherein R¹² is $\begin{array}{c} | \\ -\text{CH}- \\ | \end{array}$ or $\begin{array}{c} | \\ -\text{C}(\text{OH})- \\ | \end{array}$;


Q is a bond and R¹ is -O-CH₂-CH(OH)-;

Q is a bond and R¹ is -CH(OH)-(CH₂)₂-; or

Q is a bond and R¹ is -CH(OH)-CH₂-S(O)₀₋₂-.

30 3. A compound of any of claims 1 or 2 wherein G and G¹ are
independently selected from the group consisting of H,

-40-




wherein:

- 5 R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are independently selected from the group consisting of H, (C₁-C₆)alkyl, benzyl and acetyl;

R^3 , R^{3a} , R^4 and R^{4a} are selected from the group consisting of H, (C₁-C₆)alkyl, benzyl and acetyl; and

- 10 R , R^a and R^b are independently selected from the group consisting of H, -OH, halogeno, -NH₂, azido, (C₁-C₆)alkoxy(C₁-C₆)alkoxy and -W-R³⁰, wherein W is -O-C(O)- or -O-C(O)-NR³¹-, R³¹ is H and R³⁰ is (C₁-C₆)alkyl, -C(O)-(C₁-C₄)alkoxy-(C₁-C₆)alkyl, T, T-(C₁-C₆)alkyl, or T or T-(C₁-C₆)alkyl wherein T is substituted by one or two halogeno or (C₁-C₆)alkyl groups.

- 15 4. A compound of any of claims 1 to 3 wherein R^{26} is H or OH and G is selected from the group consisting of



wherein R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are independently selected from the group consisting of H, (C₁-C₆)alkyl, benzyl and acetyl.

20

5. A compound of claim 1 selected from the group consisting of 2,3,4-tri-O-acetyl-1-O-[4-[trans-(3R,4S)-3-[3-[(S)-acetyloxy-3-(4-fluorophenyl)propyl-1-(4-fluorophenyl)-2-oxo-4-azetidinyl]phenyl]-Beta-D-glucopyranuronic acid methyl ester;

1-O-[4-[trans-(3R,4S)-1-(4-fluorophenyl)-2-oxo-3-[3-[(S)-hydroxy-4-fluorophenyl]propyl]]-4-azetidiny]phenyl]-Beta-D-glucuronic acid;

1-O-[4-[trans-(3R,4S)-1-(4-iodophenyl)-2-oxo-3-[3-[(S)-hydroxy-4-fluorophenyl]propyl]]-4-azetidiny]phenyl]-Beta-D-glucuronic acid;

5 2,3,6-tri-O-acetyl-4-O-(2,3,4,6-tetra-O-acetyl-B-D-glucopyranosyl)-1-O-[4-[trans-(3R,4S)-3-[3(S)-acetyloxy-3-(4-fluorophenyl)propyl]-1-(4-fluorophenyl)-2-oxo-4-azetidiny]phenyl]-Beta-D-glucopyran;

1-O-[4-[trans-(3R,4S)-1-(4-fluorophenyl)-2-oxo-3-[3-[(S)-hydroxy-4-fluorophenyl]propyl]]-4-azetidiny]phenyl]-3-O-(Beta-D-glucopyranosyl)-
10 Beta-D-glucopyranose;

2,3,4,5-tetra-O-acetyl-1-O-[4-[trans-(3R,4S)-3-[3(S)-acetyloxy-3-(4-fluorophenyl)propyl]-1-(4-fluorophenyl)-2-oxo-4-azetidiny]phenyl]-Beta-D-glucopyran;

1-O-[4-[trans-(3R,4S)-3-[3(S)-hydroxy-3-(4-fluorophenyl)propyl]-1-(4-fluorophenyl)-2-oxo-4-azetidiny]phenyl]-Beta-D-glucopyranose;

1-O-[4-[trans-(3R,4S)-1-(4-fluorophenyl)-2-oxo-3-[3-[(S)-hydroxy-4-fluorophenyl]propyl]]-4-azetidiny]phenyl]-Beta-D-glucuronic acid methyl ester;

1-O-[4-[trans-(3R,4S)-1-(4-methoxyphenyl)-2-oxo-3-(3-phenyl)propyl]-4-azetidiny]phenyl]-Beta-D-glucuronic acid methyl ester;

1-O-[4-[trans-(3R,4S)-1-(4-(benzoyl)phenyl)-2-oxo-3-(3-phenyl)propyl]-4-azetidiny]phenyl]-Beta-D-glucuronic acid methyl ester;

1-O-[4-[trans-(3R,4S)-1-(4-methoxyphenyl)-2-oxo-3-(3-phenyl)propyl]-4-azetidiny]phenyl]-Beta-D-glucopyranose;

1-O-[4-[trans-(3R,4S)-1-(4-methoxyphenyl)-2-oxo-3-(3-phenyl)propyl]-4-azetidiny]phenyl]-Beta-D-glucuronic acid;

1-methyl-6-O-[4-[trans-(3R,4S)-1-(4-methoxyphenyl)-2-oxo-3-(3-phenyl)propyl]-4-azetidiny]phenyl]-Alpha-D-glucopyranoside;

1-O-[4-[trans-(3R,4S)-1-(4-(benzoyl)phenyl)-2-oxo-3-(3-phenyl)propyl]-4-azetidiny]phenyl]-Beta-D-glucuronic acid; and

1-O-[4-[trans-(3R,4S)-1-(4-fluorophenyl)-2-oxo-3-[3-[(S)-hydroxy-4-iodophenyl]propyl]]-4-azetidiny]phenyl]-Beta-D-glucuronic acid.

6. A pharmaceutical composition for the treatment or prevention of atherosclerosis, or for the reduction of cholesterol levels, comprising a compound as defined in any of claims 1 to 5, alone or in combination

-42-

with a cholesterol biosynthesis inhibitor, and a pharmaceutically acceptable carrier.

7. A process for the preparation of a pharmaceutical composition as defined in claim 6 comprising admixing a compound as claimed in any of claims 1 to 5, alone or in combination with a cholesterol biosynthesis inhibitor, with a pharmaceutically acceptable carrier.
8. The use of a compound as claimed in any of claim 1 to 5, alone or in combination with a cholesterol biosynthesis inhibitor, for the manufacture of a medicament for the treatment or prevention of atherosclerosis or for the reduction of cholesterol levels.
9. A kit comprising in separate containers in a single package pharmaceutical compositions for use in combination to treat or prevent atherosclerosis or to reduce cholesterol levels which comprises in one container an effective amount of a cholesterol biosynthesis inhibitor in a pharmaceutically acceptable carrier, and in a second container, an effective amount of a compound of any of claims 1 to 5 in a pharmaceutically acceptable carrier.
10. A method of treating or preventing atherosclerosis or reducing cholesterol levels comprising administering to a mammal in need of such treatment an effective amount of a compound claimed in any of claims 1 to 5, alone or in combination with a cholesterol biosynthesis inhibitor, wherein the cholesterol biosynthesis inhibitor, when the combination is administered, can be administered simultaneously or sequentially with the compound of claims 1 to 5.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/16823

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C07H3/04 C07H17/02 A61K31/70

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 IPC 6 C07H A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 627 418 A (BRISTOL-MYERS SQUIBB COMPANY) 7 December 1994 see the whole document ---	1
A	JOURNAL OF ORGANIC CHEMISTRY, vol. 59, no. 17, 1994, EASTON US, pages 4714-4716, XP000644310 B.K.BANIK ET AL.: "Stereospecific Glycosylation via Ferrier Rearrangement for Optical Resolution" see abstract ---	1
A	WO 95 08532 A (SCHERING CORPORATION) 30 March 1995 cited in the application see abstract --- -/--	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

11 February 1997

Date of mailing of the international search report

05 -03- 1997

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+ 31-70) 340-3016

Authorized officer

Scott, J

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/16823

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94 17038 A (SCHERING CORPORATION) 4 August 1994 cited in the application see abstract ---	1
A	CIRCULATION, vol. 80, no. 5, 1989, pages 1101-1114, XP000617211 J.L.WITZTUM: "Current Approaches to Drug Therapy for the Hypercholesterolemic Patient." cited in the application see the whole document ---	1
A	DRUGS, vol. 36, no. 3, 1988, pages 63-71, XP000617212 D.R.ILLINGWORTH: "An Overview of Lipid-Lowering Drugs." cited in the application see the whole document -----	1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/16823

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 10
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 10
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JS 96/16823

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-627418	07-12-94	US-A- 5412092	02-05-95
		AU-A- 6061194	27-10-94
		CA-A- 2120221	24-10-94
		CN-A- 1097415	18-01-95
		CZ-A- 9400966	15-12-94
		FI-A- 941894	24-10-94
		HU-A- 69373	28-09-95
		JP-A- 7002763	06-01-95
		NO-A, B 941397	24-10-94
		NZ-A- 260322	26-09-95
		ZA-A- 9402069	28-10-94
WO-A-9508532	30-03-95	AU-A- 7795294	10-04-95
		CN-A- 1131416	18-09-96
		CZ-A- 9600839	14-08-96
		EP-A- 0720599	10-07-96
		FI-A- 961300	21-03-96
		HU-A- 73852	30-09-96
		JP-T- 8509989	22-10-96
		NO-A- 961133	20-03-96
		PL-A- 313589	08-07-96
		ZA-A- 9407086	14-03-95
WO-A-9417038	04-08-94	AU-A- 6087294	15-08-94
		CN-A- 1118163	06-03-96
		CZ-A- 9501801	17-01-96
		EP-A- 0681569	15-11-95
		FI-A- 953497	20-07-95
		HU-A- 72592	28-05-96
		JP-T- 8501110	06-02-96
		LT-A, B 1764	27-02-95
		NO-A- 952884	20-09-95
		PL-A- 309978	13-11-95
		SI-A- 9400022	28-02-95
		SK-A- 91195	06-12-95
		ZA-A- 9400386	19-07-94